



APPLICATION  
FOR  
UNITED STATES LETTERS PATENT  
BY  
JOHN B. HARLEY  
JUDITH A. JAMES

---

SPECIFICATION

OF

DIAGNOSTICS AND THERAPY OF EPSTEIN-BARR VIRUS  
IN AUTOIMMUNE DISORDERS

26270-05213/80

9

\$ 154,400 101 A  
08/781296

DIAGNOSTICS AND THERAPY OF EPSTEIN-BARR VIRUS  
IN AUTOIMMUNE DISORDERS

U.S. GOVERNMENT RIGHTS

The U.S. Federal Government has rights in this invention by virtue of Grant NO. RO1 AR42460 to John B. Harley and KO8 AR01981 to Judith A. James from the National Institutes of Health.

PRIORITY INFORMATION

This application claims the benefit of U.S. Serial No. 60/019,053 entitled "EBV Vaccine for Treatment or Prevention of SLE" filed May 16, 1996 by John B. Harley, and is a continuation-in-part of U.S. Serial No. 08/160,604 filed November 30, 1993 entitled "Peptide Induction of Autoimmunity and and Clinical Symptomology" by John B. Harley, Judith A. James, and R. Hal Scofield.

Background of the Invention

This is in the area of the prevention, diagnosis, and treatment of autoimmune diseases having Epstein-Barr virus as an etiological agent.

Epstein-Barr virus infects B cells and induces a large number of different autoantibodies in the early phase of infection. The B cell proliferation and autoantibody production is eventually brought under control in nearly everyone by virus specific T cells. Thereafter, the virus remains latent, surviving in the host for the remainder of the natural life. Once the host is infected the virus continues to "reactivate" at a low level. Evidence for this reactivation is the shedding of virus in the oral cavity, infection through exchange of oral secretions, the spontaneous *in vitro* outgrowth of transformed B cells, and the spontaneous production of Epstein-Barr virus *in vitro*. The continuous presence of virus presents a significant challenge to the immune system and requires that the immune mechanisms sustain viral suppression over the many decades of remaining life. If Epstein-Barr virus causes autoimmune disease, then this feature, the sustained presence of a low level of virus in the host continuously emerging from latency, is likely to be

important in diseases that appear long after the original infection by Epstein-Barr virus.

Epstein-Barr virus is a herpes virus and is also called Human Herpes Virus 4. This virus is from the genus *Lymphocryptovirus* and subfamily *gammaherpesvirinae*. This is the only gamma herpes virus known in man. There are several very good reviews of the biology and structure of Epstein-Barr virus. The reader is referred to classic reviews (Kieff, E. and Liebowitz, D.: Epstein-Barr virus and its replication. *In* Virology, 2nd ed. Fields et al., eds. pp 1889-1921 (Raven Press, New York 1990); Miller, G.: Epstein-Barr virus. *ibid.* pp. 1921-1958; Evans, A.S. and Niederman, J.C.: Epstein-Barr virus. *In* Viral Infections in Humans, 3rd ed. Evans, A.S. ed. pp 265-292 (Plenum, New York City 1989)). Like the other herpes viruses, this is a DNA virus and has a strong propensity for latency. Once latent this virus emerges from latency at a low level throughout life. Epstein-Barr virus induces lymphoma in some non-human primates. In man Epstein-Barr virus appears to be responsible for at least infectious mononucleosis, Burkitt's lymphoma and nasopharyngeal carcinoma.

Epstein-Barr virus infects the epithelium of the upper airway, B cells and a few T cells. On B cells the viral receptor is the Complement Receptor, Type 2, also known as the CR2 receptor. Infected B cells are able to present antigen, though the virus has recently been found to produce inhibitors of antigen processing (Levitskaya, J. *Nature* 375:685-688 (1995)), and to synthesize a molecule similar to IL-10 which has profound local effects (Suzuki, T. et al *J. Exp. Med.* 182:477-486 (1995)). Depending upon what genes are expressed, latently infected B cells may not respond to stimuli in the usual way and may not provide the signals, either qualitatively or quantitatively, that would otherwise be provided. Such aberrant influences upon the normal immune response may provide the basis for subsequent autoimmune disease in some people.

Epstein-Barr virus has been known for more than three decades. For the specific example of an autoimmune disease used

herein to illustrate the principles of the invention, many others have considered a relationship between systemic lupus erythematosus and Epstein-Barr virus. The size of the separate literatures concerning lupus, on the one hand, and Epstein-Barr virus on the other are too vast to comprehensively review here. Nevertheless, over 25 years ago antibody titers were noted to be elevated against a number of viruses including rubella, measles, and parainfluenza 1 (Hollinger, F.B. et al. Bact. Proc. 131:174 (1970); Phillips, P.E. and Christian, C.L. Science 168:982-4 (1970); Hurd, E.R. et al. Arthritis Rheum. 13:724-33 (1970)). Perhaps, Dalldorf and colleagues were the first to report an evaluation of the titers of antibody against Epstein-Barr virus in lupus patients; they found no difference between lupus patients and normal controls (Dalldorf, G. et al. J. Am. Med. Assn. 208:1365-8 (1969)).

In contrast, Evans and colleagues claimed to find elevated titers of anti-Epstein-Barr antibodies relative to controls (Evans, A.S., et al. Lancet 1:167-168 (1970)). This paper generated a number of responses, all of which encouraged caution in interpreting these results or address the potential artifacts which could confuse the interpretation (Newell, G.R. and Stevens, D.A. Lancet 1:652 (1971); Evans, A.S. Lancet 1:1023-4 (1971); Gergely, L. et al. Lancet 1:325-326 (1973); Evans, A.S. and Rothfield, N.F. Lancet 1:1127-1128 (1973); Phillips, P.E. et al. Lancet 1:1449 (1973)). Much of the confusion arises from the use of immunofluorescence assays for the detection of anti-Epstein-Barr virus seroconversion. This investigative activity culminated in a remarkable study in which many participants of the controversy combined their resources to develop data they interpreted to show, "the combined approach used in this study fails to provide supportive evidence that E.B. virus is a causative agent in the connective-tissue diseases" (Klippel, J.H. et al. Lancet 2:1057-1058 (1973)). They found no difference in the titer of antibodies against Epstein-Barr virus in lupus compared to controls.

5 A Japanese group found a high frequency of antibodies  
against Epstein-Barr virus Nuclear Antigens 2 and 3 in lupus  
patient sera, compared to normal controls (Kitagawa, H., Et al.  
Immunol. Lett. 17:249-252 (1988)). Another Japanese group found  
higher levels of antibody directed against a membrane antigen  
from Epstein-Barr virus in lupus (and rheumatoid arthritis) sera  
than in controls (Yokochi, T. et al. J. Rheumatol. 16:1029-1032  
(1989)). Similarly, an Australian group found a modest increase  
in antibodies against early antigens (Sculley, D.G., et al. J.  
10 Gen. Virol. 67:2253-2258 (1986)).

An Italian group has shown that the affinity purified  
antibodies from the 95-119 region of Sm D from lupus patients  
bind the Epstein-Barr virus Nuclear Antigen-1 between amino acids  
35 and 58 (Sabbatini, A., et al. Eur. J. Immunol. 23:1146-1152  
(1993)).

The most recent contribution to this question uses both  
molecular methods to detect Epstein-Barr DNA and serologic  
methods to detect antibodies (Tsai, Y. et al. Int. Arch. Allergy  
Immunol. 106:235-240 (1995)). This study also shows no  
significant differences between lupus patients and controls.

Other diseases, including both rheumatoid arthritis and  
Sjogren's syndrome, have been explored for a possible  
relationship to Epstein-Barr virus. Robert Fox and colleagues  
presented their conception of this area in 1992 (Fox R.I., Luppi,  
25 M. and Kang H. J. Rheumatol. 19:18-24 (1992)). The evidence  
which they conclude supports a role for Epstein-Barr virus in  
rheumatoid arthritis includes: similarity between synovial and  
viral antigens, higher levels of antibodies against the Epstein-  
Barr virus Nuclear Antigens 1 and 3, and the lower ability of  
30 lymphocytes to prevent the outgrowth of autologous, Epstein-Barr  
virus infected lymphocytes (Fox, R.I. Current Opin. Rheum. 7:409-  
416 (1995)). Others have found a small increase in the frequency  
of latency for Epstein-Barr virus in rheumatoid arthritis, but a  
much larger effect for Human Herpes virus-6 (Newkirk, M.M. et al.  
35 Br. J. Rheum. 33:317-322 (1994)).

In Sjogren's syndrome Fox and colleagues note the higher level and frequency of Epstein-Barr virus in salivary gland epithelium and gland tissue (Fox, R.I. et al. J. Immunol 137:3162-3168 (1986)). Other viruses have also been considered by this author (Fox, R.I. Current Opin. Rheum. 7:409-416 (1995)).

Others have developed interesting data from Sjogren's syndrome. Pflugfelder and colleagues found evidence for Epstein-Barr virus in 80% of the lacrimal gland specimens from Sjogren's syndrome patients and in none of the controls (Pflugfelder, S.A. et al Ophthalmology 97:976-984 (1990); and Pflugfelder, S.A. et al. Am. J. Pathol. 143:49-64 (1993)). Karameris and colleagues found higher levels of hybridization between an Epstein-Barr virus DNA probe and the nuclei of salivary gland epithelial cells in Sjogren's syndrome than in controls (Karameris, A. et al. Clin. Exp. Rheum. 10:327-332 (1992)).

Others, however, found no such relationship and concluded that the frequency of Epstein-Barr virus DNA in salivary biopsy specimens was no different in patients with Sjogren's syndrome when compared with normals (Venables, P.J.W., et al. Clin. Exp. Immunol. 75:359-364 (1989); Venables, P.J.W., et al. J. Autoimmunity 2:439-438 (1989); Deacon, L.M., et al. Am J. Med. 92:453-454 (1992)). The data collected by Venables and colleagues were interpreted to show that there was "no evidence that the Epstein-Barr virus infection load is increased... [in Sjogren's syndrome]" (Venables, P.J.W. et al. Clin. Exp. Immunol. 75:359-364 (1989)), which is similar to the results of Maitland (Maitland, N.J. Am. J. Med. 96:97 (1994)). Venables and colleagues also refuted there being any abnormality in the serologic response of Sjogren's syndrome patients to Epstein-Barr virus (Deacon, E.M., et al. J. Pathol. 163:351-360 (1991)), citing their data as well as the negative serologic results of Mariette and colleagues (Mariette, X., et al. Am. J. Med. 90:286-294 (1991)).

A Japanese group found an increase in the Epstein-Barr virus production by B cells in patients with Sjogren's syndrome (Tateishi, M. et al. Arthritis Rheum. 36:827-835 (1993)).

Also, Inoue and colleagues found a minor increase in antibody levels against Epstein-Barr virus Nuclear Antigen-2 domains in Sjogren's syndrome compared to controls (Inoue, N. et al. J. Infect. Dis. 164;22-28 (1991)). Another Japanese group reported a modest elevation of anti-Epstein-Barr Nuclear antigen, anti-  
5 Early Antigen and anti-Epstein-Barr virus Viral Capsid Antigen (all measured by immunofluorescence) (Toda, I., et al. Sjogren's syndrome (SS) and Epstein-Barr virus (EBV) reactivation. In  
Lacrimal Gland, Tear Film, and Dry Eye Syndrome. D.A. Sullivan,  
10 ed. pp 647-650 (Plenum Press, New York 1994).

Nevertheless, Whittingham has proposed that Epstein-Barr virus is an etiologic agent for Sjogren's syndrome (Whittingham, S., et al. Med. Hypothesis 22:373-386 (1987)). She and her  
colleagues imagine that the Epstein-Barr viral RNAs called EBER 1 and EBER 2, which are known to bind the La autoantigen,  
facilitate overcoming tolerance to La and generating  
autoimmunity. They postulate that the combined effect of  
Epstein-Barr virus infection and autoimmunity leads to Sjogren's  
syndrome.

Morshed and colleagues published data showing an increased level of Epstein-Barr virus DNA in patients with primary biliary  
cirrhosis compared to controls from peripheral blood mononuclear cells, saliva, and fixed liver tissue (Morshed, S.A. et al. Gastroenterol. Jpn. 27:751-758 (1992)). The nuclear dot antigen  
25 is an autoantigen bound by autoantibody found in a few sera from patients with primary biliary cirrhosis. This autoantibody is also uncommonly found in lupus and rheumatoid arthritis sera. Analysis of the epitopes of the nuclear dot antigen has revealed two epitopes which have homology with Epstein-Barr virus protein  
30 sequences (Xie, K. and Snyder, M. Proc. Natl. Acad. Sci. 92:1639-1643 (1995)).

An example of double infection with Epstein-Barr virus and another virus is found in a cell line isolated from a patient with apparent multiple sclerosis (Haahr, S. et al. Ann. N. Y. Acad. Sci. 724:148-156 (1996)). The increased prevalence of  
35 seroconversion among multiple sclerosis patients, relative to

controls, has led to the suggestion that Epstein-Barr virus may be an etiologic agent in multiple sclerosis (Sumaya, C.V. et al. Ann. Neurol. 17:371-377 (1985); Bray, P.F., et al. Arch. Neurol. 40:406-408 (1983); Larsen, P.D., et al. Neurology 35:435-438 (1985); Warner, H.B. and Carp, R.I. Med. Hypothesis 25:93-97 (1988); Bray, P.F. et al. Neurology (1992)).

Because of evidence implicating Epstein-Barr virus in infectious mononucleosis, B cell lymphoma (in immunocompromised hosts), burkitt's lymphoma, nasopharyngeal carcinoma, and some cases of Hodgkin's lymphoma, there has been some activity building toward a vaccine against Epstein-Barr virus (Morgan, A.J., et al. J. Med. Virol. 29:74-78 (1989); and Morgan, A.J. Vaccine 10:563-571 (1992)). Recombinant vectors expressing gp340/220 in a bovine papillomavirus vector or in an adenovirus vector protected five of six cottontop tamarins from lymphomas that otherwise occur after infection with Epstein-Barr virus (Finerty, S., et al. J. Gen. Virol. 73:449-453 (1992)). A subunit of the gp340/200 in alum protected three of five cotton top tamarins from lymphomas (Finerty, S., et al. Vaccine 12:1180-1184 (1994)), suggesting that this strategy might not be especially effective. A trial of an Epstein-Barr virus vaccine of gp340/220 in a Vaccinia virus vector has been reported from China and failed to protect a third of those immunized (Gu, S. et al. Dev. Biol. Stand. 84:171-177 (1995)).

A variety of therapies have been attempted against Epstein-Barr virus. These include inducing the lysis cycle in cells latently infected by virus (Gutierrez, M.I., et al. Cancer Res. 56:969-972 (1996)). Patients with the Epstein-Barr virus related lymphomatoid granulomatosis have been treated with interferon-alpha 2b with the preliminary impression that the treatment was successful (Wilson, W.H., et al. Blood 87:4531-4537 (1996)). Cycloheximide has been demonstrated to be useful in vitro (Ishii, H.H., et al. Immunol. Cell Biol. 73:463-468 (1995)). Therapy with a T cell line has been attempted (Kimura, H. et al. Clin. Exp. Immunol. 103:192-298 (1996)), as has adoptive transfer of gene-modified virus-specific T lymphocytes (Heslop, H.E. et al.



5 Nature Med. 2:551-555 (1996)). Data available do not appear to particularly support the use of acyclovir for Epstein-Barr virus infections (Wagstaff, A.J., et al. Drugs 47:153-205 (1994)), though FK506 (a relative of cyclosporine) may have some benefit (Singh, N., et al. Digestive Dis. Sci. 39:15-18 (1994)). Monoclonal antibodies have been used to treat the virus-induced lymphoproliferative syndrome (Lazarovots, A.I., et al. Clin. Invest. Med. 17:621-625 (1994)).

10 It is therefore an object of the present invention to provide strategies to prevent autoimmune disease by vaccination with vaccines based upon Epstein-Barr virus or upon the structure of Epstein-Barr virus.

It is a further object of this invention to provide vaccines based upon Epstein-Barr virus or upon the structure of Epstein-Barr virus which will have little risk of inducing autoimmune disease.

It is a further object of this invention to provide diagnostics which will identify people exposed to Epstein-Barr virus who are at an increased risk for autoimmune disease and, alternatively, those who are at decreased risk for developing autoimmune disease.

It is a further object of this invention to provide for the application of antiviral therapy directed against Epstein-Barr virus in the treatment of autoimmune disease.

25 It is a further object of this invention to provide diagnostics and therapeutics for autoimmune disease based upon the changes induced by the host by Epstein-Barr virus.

### Summary of the Invention

30 Data consistent with autoimmune disease being caused by Epstein-Barr virus are shown. Some of the features of the mechanism in the specific example of the anti-Sm autoantibody response were found in systemic lupus erythematosus. Based on this evidence, an effective vaccine would prevent the autoimmune disease in those vaccinated, modified or administered so that the vaccine is not itself capable of inducing autoimmune disease. In

35

the case of anti-Sm, structures to be avoided in an Epstein-Barr virus-derived vaccine have been identified.

Differences have been identified in the immune responses to Epstein-Barr infection between individuals who develop a specific autoimmune disease and those who do not. These differences are used to distinguish those who are at greater risk for developing specific autoimmune diseases from those who are a lesser risk.

Assuming Epstein-Barr virus causes autoimmune disease and that Epstein-Barr virus remains latent in the patient for life, reactivation of the virus from the latent state is important in generating or maintaining the autoimmune response that culminates in autoimmune disease. Cells infected with latent virus may also encourage autoimmunity. Based on the understanding that reactivation or latency are important to produce or sustain autoimmunity, then therapies directed against Epstein-Barr virus will also be effective therapies for the autoimmune manifestations of disease for which Epstein-Barr virus is responsible.

#### Brief Description of the Drawings

Figures 1A-D, graphs of absorbance indicative of the antigenicity of overlapping octapeptides of an Sm B/B' polypeptide. Each number along the abscissa indicates the first amino acid of an octapeptide that begins with this amino acid (octapeptide number, 1-8, 2-9, 3-10, etc.). Each octapeptide overlaps its neighbor by seven amino acids. Figure 1A shows the background reactivity of the octapeptides with anti-human IgG conjugate alone. Figure 1B shows the reactivity of the octapeptides with a normal human serum. Figure 1C shows binding of a serum from a patient with systemic lupus erythematosus who precipitates the nRNP autoantigen, but not Sm autoantigen. Figure 1D demonstrates the reactivity of a representative patient who has both anti-Sm and anti-nRNP autoantibodies as determined by specific precipitin formation in Ouchterlony immunodiffusion. From J.A. James and J.B. Harley (J. Immunol. 148;2074-2079, 1992).

Figure 2. Mean binding to overlapping octapeptides by Sm and nRNP precipitin positive lupus patients showing the major epitopes (as defined as being greater than 0.5 O.D., or a mean of 10 standard deviations above the mean binding of normal sera). Binding below the threshold of 0.325 O.D., or the mean of normals plus two standard deviations, has been omitted for clarity. The standard error is depicted as an open box below the mean in the bar representing the mean of the lupus patient binding. From J.A. James and J.B. Harley (J. Immunol. 148;2074-2079, 1992).

Figure 3 is a graph of binding to selected peptides from the B/B', C and D spliceosomal proteins and from Epstein-Barr virus nuclear antigen-1 proteins (PPPGRRP, GRGRGRGG, and RGRGRER) in six lupus patient sera (which are anti-Sm and anti-nRNP precipitin positive) (black) and five control sera (shown overlaid in white).

Figures 4A-D are graphs of the development of an anti-Sm B/B' response in one anti-Sm and anti-nRNP precipitin positive patient. Binding to 211 octapeptides of Sm B/B' is presented as absorbance (X 1000) at 405 nm. Figure 4A presents an example of the background binding found from a normal human serum. Figure 4B presents the binding of the first serum available after presentation with lupus (from April, 1986) and Figure 4C and 4D present the binding of subsequent sera (from July, 1987 and December 1988, respectively). Solid arrows indicate the PPPGMRPP octapeptides and the open arrow the PPPGIRGP sequence. From J.A. James et al (J. Exp. Med. 181:453-461 (December 1995)).

Figures 5A-C are graphs of the binding of PPPGRRP-MAP<sup>TM</sup> immunized rabbit sera and a Freund's control serum to the PPPGRRP-MAP<sup>TM</sup> (Figure 5A), PPPGMRPP-MAP<sup>TM</sup> (Figure 5B), and nRNP/Sm antigen (Figure 5C) by solid phase ELISAs (enzyme-linked immunosorbent assays). The bleeds of the rabbits are along the abscissa and the absorbance (X 1000) on the ordinate. The preimmune serum (pb) and post-immunization bleeds 1 through 14 (Im1 to Im14) which span 52 weeks are indicated. The PPPGRRP peptide is found in the Epstein-Barr virus Nuclear Antigen-1 (EBNA-1).

Figures 6A-D are graphs of the binding of two rabbit sera to the overlapping octapeptides of the spliceosomal proteins B/B' (Figures 6A and 6C) and D (Figures 6B and 6D). The two rabbits (numbers 40 and 41) were immunized with PPPGRRP-MAP<sup>TM</sup> 86 days earlier. The serum in Figures 5A and 5B binds to nRNP/Sm, and is positive in the antinuclear antibody assay and the anti-double stranded DNA assay (*Crithidia lucilea* kinetoplast fluorescence assay) while the serum from the second rabbit was not positive in any of these other assays.

Figure 7 is a graph of the binding to the peptide GAGAGAGAGAGAGAGAGAGAGAGA from Epstein-Barr virus Nuclear Antigen-1 by lupus sera who all had anti-Sm and anti-nRNP precipitins, as compared to normal control sera.

Figures 8A-E are graphs of the binding to the overlapping octapeptides from Epstein-Barr virus Nuclear Antigen-1. Each octapeptide overlaps its neighbor by seven amino acids. Most of the glycine-alanine repeat has been omitted to avoid unnecessary redundancy. The binding of three controls are presented in Figures 8A, 8B and 8C and that of two lupus sera in Figures 8D and 8E. Figure 8A is from a normal who has no evidence of having been infected by Epstein-Barr virus by the assay for anti-Epstein-Barr virus Viral Capsid Antigen IgG. The other sera presented (Figures 8B through 8E) are all positive in this assay.

### Detailed Description of the Invention

In the United States, about 95% of the adult population has been, and continues to be, infected with Epstein-Barr virus. Observations described herein are consistent with a small proportion of these developing autoimmune disease, related to this virus. Other factors are also likely to be important in the development of autoimmune disease, but are not essential to understand in order to develop therapeutics and diagnostics for use in diagnosing, treating and preventing or ameliorating autoimmune diseases involving Epstein-Barr virus as the etiologic agent. Epstein-Barr virus is the probable etiologic agent for

nearly all cases of lupus, which serves as an example of autoimmune disease.

5       Diagnostics and therapeutics derived from the discovery that Epstein-Barr virus causes autoimmune disease as applied to the prevention, diagnosis and treatment of autoimmune disease are described herein. Systemic lupus erythematosus (lupus) is the particular autoimmune disease evaluated and for which data have been obtained. Within lupus, the work on a molecular understanding of the relationship between anti-Sm and systemic lupus erythematosus and the relationship of anti-Sm autoantibodies to Epstein-Barr virus is the best illustration of the data supporting these diagnostics and therapeutics.

10       The experiments described herein to address Epstein-Barr virus in lupus were guided by the results of immunochemical studies, not by the previous studies. These data pointed toward a curious mechanism in the anti-Sm autoantibody system in lupus which could involve Epstein-Barr virus.

15       The technology applied to the problem is very important in two ways. First, the assays for anti-Epstein-Barr virus antibodies have been dramatically improved. The classic method is to evaluate antibody binding to an Epstein-Barr virus infected cell line by immunofluorescence. This assay is dependent upon the expression of different Epstein-Barr virus proteins, depending upon whether the cell line is producing virus or the virus is latent. The autoantibodies of lupus often render these assays uninterpretable, making their use in lupus especially problematic.

25       Consequently, solid phase assays using enriched preparations of the surface antigen constitute a major improvement. The surface antigen is largely composed of a glycoprotein called gp340/220 or the Viral Capsid Antigen. These preparations of surface antigens have had many interfering substances removed. Epstein-Barr virus infection in man virtually always generates antibodies against this surface antigen. The assay is much simpler than is the cell line immunofluorescence assay and subject to much less variation in interpretation.

Second, molecular methods have been designed and developed to detect Epstein-Barr virus which appear to be at least as reliable as the serologic methods, and may even be superior. In normal adults Epstein-Barr virus infects only about one in every 20,000 to 500,000 B cells (Miyashita, E. M. et al. Cell 80:593-601 (1995)). B cells usually constitute only about 8% of the peripheral blood mononuclear cells. The vanishingly small quantity of Epstein-Barr DNA is lost in a relative ocean of genomic human DNA and is very difficult to detect. The improved sensitivity and specificity of detection of Epstein-Barr virus DNA improves the measurement made and leads to more accurate interpretation of the data. The older methods detected Epstein-Barr virus DNA in the peripheral blood in about 70% of individuals who had serologically converted, while the method described herein appears to detect more than 95% of those who have seroconverted and a few who have not seroconverted.

Reliable assays were used to address the prevalence of seroconversion and infection in the cases and controls in a way that took optimal advantage of the known properties of the viral infection. Others have selected sub-optimal study populations, have poorly chosen their controls, or have focused upon the quantitative level of antibody rather than qualitative evidence for infection, in addition to the technical problems outlined above in reliably detecting anti-Epstein-Barr virus seroconversion and Epstein-Barr viral DNA.

The data in the anti-Sm autoantibody system, discussed below, are used as a model in which the antigen presenting capacity of the B cell is important in generating autoimmunity. For example, the PPPGRRP structure is found in Epstein-Barr virus Nuclear Antigen-1. This sequence induced autoimmunity against the Sm B/B' of a rabbit after immunization. This autoimmunity not only included the related PPPGMRPP of Sm B/B', but also many other structures of B/B'. When the B cell generates a receptor that binds PPPGRRP and PPPGMRPP then this B cell is capable of presenting the spliceosome to the immune system. Of course, once this cross reacting autoantibody is produced, then it may

facilitate spliceosomal autoimmunity. Epstein-Barr virus is important because the immune control of the infected B cell is altered by the infection, rendering autoimmunity more likely. This mechanism can be directly extended to other antigens to generate other immune responses (both cellular and humoral) which lead to a variety of autoimmune diseases. Also, double infection with Epstein-Barr virus and another virus would extend the immune regulatory abnormalities to the antigens of the second virus.

#### Autoimmune Diseases

There are a large number of disorders in man that are thought to be autoimmune. These include systemic lupus erythematosus, autoimmune thyroid disease (Graves' disease or Hashimoto's thyroiditis), autoimmune beta islet disease of the pancreas (more commonly referred to as juvenile or Type 1 diabetes mellitus), primary biliary cirrhosis and many others. The particular disorders listed above are thought to involve antibodies produced in the host (the patient, in this instance) which bind to constituents of self. These antibodies are called autoantibodies. The particular constituent of self bound by the autoantibodies is associated with the different disorders. For example, anti-mitochondrial autoantibodies are associated with primary biliary cirrhosis. Anti-acetylcholine receptor autoantibodies are associated with myasthenia gravis. The list of such autoantibodies is quite long and often only one or a few autoantigens are bound by autoantibodies in each particular disorder. Systemic lupus erythematosus (or abbreviated as lupus herein) is an exception to this tendency, since many autoantibodies may be found in the disease and since patients do not necessarily share any particular autoantibody specificity. Anti-Sm (which is an anti-spliceosomal autoantibody specificity) is one of the autoantibodies closely associated with systemic lupus erythematosus, but even this autoantibody is found in only a minor fraction of patients with systemic lupus erythematosus. (Please refer to a review of this area (Harley, J.B. and Gaither, K.K. Rheum. Dis. Clin. N. Amer. 14:43-56 (1992)).

It is not sufficient just to produce autoantibodies. There must be some consequence of their presence in order to develop pathology which culminates in clinical disease. There are many instances of detecting autoantibodies in the absence of any detectable clinical illness. Autoantibodies may realize their pathologic potential by binding their antigen in the circulation. They then become part of circulating immune complexes. They may deposit in tissues, induce an inflammatory response, and cause tissue injury, as appears to occur in lupus. Autoantibodies may interfere with the functioning of receptors or otherwise activate cells as may happen in Wegener's granulomatosis or Graves' ophthalmopathy. Autoantibodies may simply block normal functioning of a protein, as happens to the acetylcholine receptor in myasthenia gravis. No doubt there are other mechanisms by which autoantibodies encourage clinical illness.

These mechanisms involve humoral autoimmunity; that is, autoimmunity that is mediated by autoantibodies. There is another form of autoimmunity mediated by cells, in particular T cells. Multiple sclerosis is thought by some to be an example of a disease that is mediated by autoimmune T cells. Although the methods and compositions described herein are particularly concerned with humoral autoimmunity, it is expected that cellular autoimmune processes are also involved with producing autoimmunity as a consequence of Epstein-Barr virus infection. As in many situations, one skilled in the art would expect cellular immune mechanisms to dominate in some individuals and humoral mechanisms to dominate in others. This situation would be expected to give rise to different clinical expression of disease. Tuberculous and lepromatous leprosy are examples where differences in the dominant form of the immune response lead to profound differences in the clinical illness, despite being caused by the same organism.

The traditional distinction between humoral and cellular immune mechanisms are being reevaluated under a new paradigm. T cells appear to have the capacity to respond in at least two ways. These cells are called Th1 and Th2, for T helper cells,



5 Type 1 and 2. Characteristic cytokine production profiles are often used to distinguish these different responses. Th1 responses tend to be the more traditionally appreciated cellular immune responses. Th2 responses lead, among other consequences, to more of an antibody response and are more aligned with the classic humoral response. However, these boundaries do not appear to operate strictly since some types of antibody are more likely found in Th1 responses and the Th2 response clearly has its cellular component. Most autoimmune diseases probably have important components of both humoral and cellular autoimmunity.

10 Below, systemic lupus erythematosus is used as a particular example of an autoimmune disease. Data are presented which is consistent with the position that Epstein-Barr virus causes this autoimmune disorder. Lupus is one of many autoimmune diseases that are likely to share basic features, such as the causative agent being Epstein-Barr virus.

#### Definitions:

As used herein, autoimmune diseases are diseases that are primarily autoimmune, as well as diseases which do not appear to be primarily autoimmune but have immune manifestations involving immunoglobulins, antigen specific B cell surface receptors (surface immunoglobulin), or antigen-specific T cell receptors. Examples of diseases which fall into these categories are systemic lupus erythematosus, Sjogren's syndrome, rheumatoid arthritis, juvenile onset diabetes mellitus, Wegener's granulomatosis, inflammatory bowel disease, polymyositis, dermatomyositis, multiple endocrine failure, Schmidt's syndrome, autoimmune uveitis, Addison's disease, adrenalitis, primary biliary cirrhosis, Graves' disease, thyroiditis, Hashimoto's thyroiditis, autoimmune thyroid disease, pernicious anemia, gastric atrophy, chronic hepatitis, lupoid hepatitis, atherosclerosis, presenile dementia, demyelating diseases, multiple sclerosis, subacute cutaneous lupus erythematosus, hypoparathyroidism, Dressler's syndrome, myasthenia gravis, autoimmune thrombocytopenia, idiopathic thrombocytopenic purpura, hemolytic anemia, autoimmune hemolytic anemia, pemphigus



polychondritis, malignant melanoma, cryoglobulinemia, hepatitis B virus infection, hepatitis C virus infection, Waldenstrom's macroglobulinemia, mumps virus infection, and any other disorder in which the specific recognition of the host by immunoglobulin, B cell surface receptor (surface immunoglobulin), or T cell receptor is suspected or shown to be important in any aspect of the pathogenesis of the clinical illness.

Immunization is any procedure which leads to a cellular or humoral immune response directed against an identifiable and specific antigen, usually the immunogen. An antigen is a substance that is bound by antibody. Sometimes an antigen is also referred to when meaning a substance against which any immune response is directed and that it may be bound by antibody or lead to a cellular immune response. An autoantigen means a constituent of self that binds antibody (making it an autoantibody) or that induces a cellular response, for example, by a T cell. The spliceosome is that molecular apparatus, composed of RNA and protein, which splices heteronuclear RNA, thereby removing the introns from the coding sequence of RNA. The cellular response may be assayed by presentation of a peptide from the autoantigen, proliferation, cell activation, the prevention of cell activation, secretion of cytokines, activation of apoptosis, or other indication of an effect of the presence of the autoantigen. An autoantibody is any immunoglobulin, antigen specific B cell surface receptor (surface immunoglobulin), or antigen specific T cell receptor directed against self protein. Such T cell receptors usually bind peptides which themselves are bound by histocompatibility molecules. The T cell receptor usually binds to both the peptide and the histocompatibility molecule.

Therapy is a treatment by medical or physical means. A "treatment" is the composition used for treating a condition. Antiviral therapy is the use of a treatment in an effort to suppress or eliminate a virus, for example, suppression, elimination or other amelioration of the effect of Epstein-Barr virus. Peptides are small proteins composed of amino acids

covalently bound to one another by peptide bonds. Peptides may be prepared by an *in vivo* mechanism, as in life, by using the nucleic acid encoding for the sequence of the peptide produced, or *in vitro* using peptide chemistry. A vaccine is a composition or preparation that is used in an attempt to prevent a disease or ameliorate a disease. A disease is any illness.

Seroconversion means that the subject has developed antibodies of sufficient magnitude in the serum to conclude that the subject has made an immune response against the agent or substance of interest. Usually, this is the result of immunization (vaccination) or infection. Seropositive means that there are a sufficient quantity of antibodies with sufficient affinity to conclude that seroconversion has occurred. Seronegative means that the quantity and affinity of antibodies are not sufficient to conclude that seroconversion has occurred. In this application, the terms "lupus" and "systemic lupus erythematosus" are used interchangeably.

The single amino acid code is used in the figures and following examples, as follows:

A - alanine	I - isoleucine	R - arginine
C - cysteine	K - lysine	S - serine
D - aspartic acid	L - leucine	T - threonine
E - glutamic acid	M - methionine	V - valine
G - glycine	P - proline	Y - tyrosine
H - histidine	Q - glutamine	

### Therapeutic and Diagnostic Compositions

#### Vaccines

Immunity against a viral infection can be induced using either peptides, viral proteins or other components of the virus such as carbohydrate components, substances which imitate structures of the virus, or the virus. In the preferred embodiment, the vaccine is based on the viral proteins wherein the epitopes cross-reactive with the splicesomal proteins are deleted. In other embodiments, the vaccine is based on viral proteins where epitopes cross-reactive with antibodies to other

known autoantigens are deleted or altered to decrease their immunoreactivity with autoantibodies. In still other embodiments, the vaccine includes the virus, either live, modified or inactivated, or components thereof, in a vehicle which can be administered in a dosage form and over a schedule eliciting a strong immune response to kill the virus, but which does not elicit an autoimmune response.

An unmodified vaccine may be useful in the prevention and treatment of autoimmune disease related to Epstein-Barr virus. The dose, schedule of doses and route of administration may be varied, whether oral, nasal, vaginal, rectal, extraocular, intramuscular, intracutaneous, subcutaneous, or intravenous, to avoid autoimmunity and, yet, to achieve immunity from Epstein-Barr virus infection. The response to the unmodified vaccine may be further influenced by its composition. The particular adjuvant employed (its concentration, dose and physical state), concentration of the virus in the vaccine, and treatment of the unmodified vaccine with physical environmental changes, for example, temperature and pressure, the particular buffer, and the particular preservative(s) (if any) will be selected to reduce the likelihood of developing an autoimmune disorder, for example using the animal strains discussed below. This same or different vaccine may be useful in reducing or eliminating the effect of an existing latent or active Epstein-Barr virus infection upon autoimmunity.

Peptides of up to about forty amino acids, more preferably between four and twenty-five amino acids, most preferably eight amino acids, can be synthesized using any one of the methods known to those skilled in the art. In general, an epitope of a protein is composed of between three or four and eight amino acids (see Watson et al., "Certain Properties Make Substances Antigenic," in Molecular Biology of the Gene, Fourth Edition, page 836, paragraph 3, (The Benjamin/Cummings Publishing Company, Menlo Park, 1987). As used herein, the peptides can contain the entire native epitope, or portions thereof sufficient to react with autoantibody.

Although described in the literature with reference to specific sequences encoding viral proteins or the autoantigens, a number of substitutions using natural or synthetic amino acids can be made in the peptides to yield a peptide acting as a linear epitope that is functionally equivalent to the disclosed sequences, for example, as demonstrated by James and Harley, J. Immunology 148:2074-2079 (April 1992). Accordingly, the term linear epitope as defined by a specific sequence is used herein to include peptides having substitutions yielding a peptide bound in an equivalent manner or extent by an antibody or autoantibody. For example, using monoclonal antibodies against peptide determinants of Sm B/B', substitution studies demonstrated that

A, G, and S can substitute for R in PPPGMRPP in the binding of one antibody, KSm3. Analogously, F, H, T, V and Y can substitute for I in PPPGIRGP in the binding of KSm3. (SEQ ID NO: 4)  
(SEQ ID NO: 5)

Solid phase binding of autoantibodies to peptides has proven useful for examining sequential linear epitopes, also referred to as "linear" or "sequential". These have been useful to define important residues in epitope structure. This approach may or may not be less useful in defining conformational epitopes or regions where two or more linear, but not sequential, epitopes are brought together by the tertiary structure. The particular structural relationships between the autoantigens and autoantibody with particular regard to the particular conformation assumed by the peptide determines what can be learned by this approach. In addition, although many peptides may assume conformations in solution that are not found in the native protein structure, true epitopes may still be delineated by this method. Those peptides that tend to have a structure similar to that found in the native molecule are expected to usually be bound by a larger proportion of the autoantibodies that bind the analogous sequence on the native protein and/or may be bound with greater affinity.

It is believed that naturally arising human lupus follows a progression similar to that induced in the rabbit model described in the examples. Using this model, an immune response to a

peptide, one very similar to a region of EBNA-1, is the seminal, initiating event for the subsequent autoimmunity and disease manifestations of lupus, where they occur. A structure, such as a peptide, that is capable of inducing autoimmunity is not necessarily identical to the structure found in the autoantigen. Indeed, it is possible that these structures would commonly be at least slightly different, when comparing the substance that induces the autoimmune response and the analogous structure in the autoantigen. On the other hand, there must be a basis for the non-autoantigen substance to induce autoimmunity. This is best identified as a cross-reaction wherein the immune recognition molecule binds, though not necessarily equally, to both the non-autoantigen substance as well as to the autoantigen.

The proposed mechanism is as follows. An immune response against a non-autoantigenic substance occurs. Some fraction of the antibodies thereby produced recognize the autoantigen and hence are autoantibodies. These autoantibodies facilitate the processing and antigen presentation of the autoantigen via the B cell surface immunoglobulin which serve as receptors, in this case for autoantigen, or via immunoglobulin cell surface receptors that are found on a variety of cells capable of antigen presentation. Once this occurs the immune response expands to other structures of the autoantigen and a full-blown, complete autoimmune response against the autoantigen ensues, which can result in clinical illness.

Based on this mechanism, the autoimmune responses progress from one or a few initial antigenic structure(s) to a much more complex response focused upon the autoantigen. Elucidating the pattern of progression and understanding the relationship of autoimmune serologic findings to clinical manifestations places the physician in a strong position to accurately prognosticate and prepare patients and their families for the more likely outcomes.

In the case of systemic lupus erythematosus and the anti-Sm response, one method is to repeat the assays determining autoantibody binding to peptides over time. The effect of the

peptide *in vitro* on cells from patients can also be measured. Proliferation, secretion of cytokines, interferons and other substances, expression of cell surface molecules and activation are typically useful diagnostic indicators.

5 This strategy to generate autoimmunity can also be used to develop reagents that are useful in diagnosis or treatment of autoimmune disorders. Animal antibodies that compete with or otherwise facilitate the identification of particular fine specificities of binding can be important in evaluating  
10 prognosis. Moreover, the peptide binding pattern to the octapeptides from the nRNP A protein show two different patterns. It should be possible to correlate a particular pattern found in a patient to obtain an indication of the stage the disease currently is at as well as the clinical prognosis. Reagents developed as a consequence of immunizing animals with autoantigenic peptides could be used to identify these differences. Such reagents include antisera, T cell lines, subsets of antibodies, individual antibodies, subsets of cells bearing a subset of the T cell receptors, individual T cell  
15 receptors, and cytokines and other substances elaborated by cells from the animal. The antibodies and T cell receptors are construed to include recombinant antibodies or T cell receptors derived from a peptide-immunized animal.

25 The RNA-protein particles which are the major autoantigens may now be purchased commercially. The reagents made available by the animal model of autoimmunity described herein will be useful in the manufacturing and testing of autoantigens. Affinity purification using animal antiserum (absorbed or otherwise prepared) could be used for purification of the  
30 naturally occurring autoantigens.

Having a mechanism of disease provides the opportunity to apply new strategies for prevention of disease and for specific immunologic correction of the immune abnormalities that lead to disease, and therefore more accurately design the therapy.

35 For example, with the realization that the generation of autoimmune disease can be divided into phases comes the



B  
5 D appreciation that the therapeutic opportunities will be similarly partitioned. As a specific example, the influence of vaccination with an analog of PPPGMRPP will be different depending not only upon the structure of the immunogen, but also upon the pharmaceutical carrier, upon the maturity of the autoimmune response against PPPGMRPP and Sm, and upon other therapeutics that may be administered concomitantly. Such therapeutics include drugs as well as biologics, such as cytokines, immunoglobulins, and interferons, among others.

10 D As a second specific example, mice strains exist which produce lupus autoimmunity, as well as strains which do not produce lupus autoimmunity, after immunization with PPPGMRPP-MAP™. The genetic, biochemical, and physiological differences between these strains can be used to develop diagnostics, including genetic markers and risk factors, and therapeutics for autoimmune disease, especially lupus, using techniques known to those skilled in the art.

B  
15 00131296-011397 As discussed above, based on the data confirming that Epstein-Barr virus is an inducing agent for lupus, it is possible to design therapeutics to prevent or inhibit further progression of lupus by vaccinating with Epstein-Barr virus, or components thereof, using standard vaccination procedures, most preferably after alteration or removal or masking of the sites which elicit the autoimmunity.

25 The peptides can be used therapeutically in combination with a pharmaceutically acceptable carrier. The peptides can be administered in a dosage effective to block autoantibodies or as a vaccine to block the production of autoantibodies, by eliciting a protective immune response against non-autoantigenic regions of the pathogen. The peptide acts as a functional antagonist by binding to antibody that does not stimulate or activate the immune cells and thereby block the immune response to the autoantigens.

30 35 Pharmaceutical carriers are known to those skilled in the art. These most typically would be standard carriers for administration of vaccines to humans, including solutions such as

sterile water, saline, and buffered solutions at physiological pH. Peptides used as vaccines are most preferably administered intramuscularly or subcutaneously. Other compounds will be administered according to standard procedures used by those skilled in the art.

Alternatively, the peptides used for treatment might include peptides homologous to an identified antigenic sequence. These peptides, either free or bound to a carrier, could be delivered to a patient in order to decrease the amount of circulating antibody with a particular specificity. In addition, knowledge of the cross-reacting epitopes between a foreign antigen and an autoantigen may allow for re-induction of tolerance. It is well known in experimental models of the immune response that the response can be suppressed and tolerance induced by treatment with the antigen. Peptide therapy with the cross-reacting sequences may be a useful therapy in autoimmune diseases.

The amino acid sequences can also be used to make agents for neutralizing circulating antibodies or immobilized on substrates in extracorporeal devices for specific removal of autoantibodies, using methodology known to those skilled in the art.

#### *Diagnostics*

Individuals who are not at as great a risk for developing autoimmune disease can be identified by reactivity to the various peptides, for example, as demonstrated in the examples where individuals who are not prone to develop lupus are characterized by antibodies to GAGAGAGAGAGAGAGAGAGAGAGA. Other structures derived from Epstein-Barr virus can be used to predict who will develop autoimmune disease. For example, GAGAGAGAGAGAGAGAGAGAGAGA is a commonly identified antigen from Epstein-Barr virus in normal individuals, while lupus patients do not tend to bind this structure. An individual identified at risk for the development of an autoimmune disease, but who does not yet manifest autoimmunity or symptoms of the disease, may require a special therapeutic approach. This is an opportunity to induce immune suppression before the process leading to autoimmune disease is initiated. Strategies such as intravenous

administration of large amount of the initiating structure is known to induce tolerance. Small sub-immunogenic doses of the initial immunogen can also be used to induce tolerance.

5 There is a limited opportunity to interrupt or redirect an immune response that has been initiated against the first components of the autoantigen. Here again the induction of suppression by the use of the component peptides or analogs thereof with or without concomitant drugs or biologics has the potential to inhibit progression into an autoimmune disorder. 10 Once autoimmunity against the autoantigen is established, the use of component peptides or their analogs with or without concomitant drugs or biologics may interrupt the course of the autoimmune response, thereby ameliorating the illness.

25 The animal model provides an opportunity to optimize ways of interrupting and reversing the autoimmune process. For example, it has been observed that one of the rabbits immunized with Map-PPPGMRPP (SEQ ID NO: 4) seemed to improve clinically somewhat after developing the most severe manifestations of systemic autoimmunity. If this result is the effect of a particular antibody, then this antibody may have the capacity to influence the maturation of the immune response toward alleviating the disease in other species. For example, such an antibody could be isolated by biochemical methods, by recombinant DNA methods or by hybridoma monoclonal methods, humanized using standard technology and then administered to patients as a specific therapeutic agent for disease. T cell receptors or cytokines could be equally useful.

#### Assays

30 Subsets of antigenic peptides have the potential to identify patients at risk for particular clinical manifestations or patients in particular prognostic groups. The peptides can be used in combination in assays, such as the solid phase assay, to classify patients.

35 Specifically, the peptides that are bound by autoantibodies in patients characterized by specific disorders, such as renal disease or central nervous system involvement, are selected and combined in an assay, such as an ELISA for a test to detect the

collection autoantibodies that bind this particular collection of peptides. Using a mixture of peptides may increase the efficiency and reliability of such assays, as compared with using a single autoantigen, or a single peptide.

5 The peptides can be used in solution or immobilized to a solid substrate, such as a gel suitable for affinity chromatography, or a multi-well plate, using standard techniques such as the commercially available cyanogen bromide.

The present invention will be further understood by reference to the following non-limiting examples.

10 **Example 1: The relationship of anti-Sm to Epstein-Barr virus.**

25 The anti-Sm and anti-nuclear ribonucleoprotein (nRNP) autoantibody specificities have an Ouchterlony precipitin reaction of partial identity. This is because the epitopes bound are found on two antigens which share some, but not all antigenic features. The Sm autoantigen may be found in association with the U1, U2, U5 or U4/6 spliceosomal particles. These are composed of specific U RNA and particular peptides. The nRNP autoantigen is composed of only U1 RNA and particular unique peptides. The anti-Sm antibodies bind to the B/B' or D spliceosomal peptides in Western blot while the anti-nRNP autoantibodies bind 70K, A or C peptides (Lerner, M.R. and Steitz, J.A. Proc. Natl. Acad. Sci. USA 76:5495-5499 (1979); Hinterberger, M. J. Biol. Chem. 258:2604-2613 (1983); and Petterson, I. et al. J. Biol. Chem. 259:5907-5914 (1984)). (B/B' and D are found in the U1, U2, U5 and U4/6 spliceosomal particles. A protein termed the N protein or the Sm N protein is closely related to B and B'. 70K, A and C are usually only found in U1.) Anti-Sm precipitins are found in about 10% to 20% of lupus patients. Anti-Sm is sufficiently specific for this diagnosis that its presence in the serum has become part of the 30 1982 Criteria for the Classification of Systemic Lupus Erythematosus (Tan E.M., et al Arthritis Rheum. 25:1271-1277 (1982)).

35 *Antigenicity of B/B' peptides assessed by measuring antibodies binding to octapeptides synthesized in vitro from the sequence of the B/B' Sm peptide.*

The autoantigenicity of Ro (also known as the SS-A) autoantigen was reported by Scofield, R.H. et al. Proc. Natl. Acad. Sci. USA 81:3343-3347 (1991). The antigenicity of the B/B' peptides was assessed by measuring antibodies binding to octapeptides synthesized *in vitro* from the sequence of the B/B' Sm peptide, as described by James, J.A. and Harley, J.B. J. Immunol. 148:2074-2079 (1992). Representative data from an anti-Sm precipitin positive patient are shown in Figure 1D and controls in Figures A-C. Note that eleven groups of peptides are bound by the sera containing anti-Sm autoantibody. One of the most unexpected observation was that all of the anti-Sm sera tested bound almost the identical octapeptide structures (Figure 2).

The sequence PPPGMRPP is repeated three times in Sm B' (Van Dam, A. et al. EMBO J. 8:3853-60 (1989)). The N protein, which is very homologous to the B/B' protein, has three repeats of PPPGMRPP and a closely related sequence PPPGIRGP is also found once (Schmauss C. et al. Nuc. Acid Res. 17:1733-43 (1989)). PPPGIRGP is not found in the B' protein; rather, PPPGMRGP is found in its place. Much confusion has surrounded this area due to the original publication of the Sm B/B' sequence (Rokeach L.A. et al. J. Biol. Chem. 264:5024-30 (1989); Sharpe N.G. et al. FEBS Lett. 250:585-90 (1989)) which was later found to actually produce what is now termed the N protein or the Sm N protein (Schmauss C. et al. Nuc. Acid Res. 17:1733-43 (1989)). Lupus patients with anti-Sm precipitins all have antibodies which bind to both Sm B (a truncated version of Sm B'), Sm B' and Sm N. Substantial cross-reactivity occurs between the three proteins.

Such repeated structures are sometimes particularly immunogenic. These anti-Sm sera also bind to some of the known sequences which are structurally similar to PPPGMRPP (Figure 3). These sequences included PPPGRRP, which is found in the Epstein-Barr Nuclear Antigen-1 (EBNA-1) protein. GRGRGRGG and RGRGREK are also sequences from Epstein-Barr virus Nuclear Antigen-1, but these are similar to a major antigenic epitope of Sm D in lupus patients, GRGRGRGRGRGRGRGRGRGRGGPRRR (James, J.A., et al. Clin.

Exp. Immunol. 98:41-426 (1994)). (See GenBank accession code: gb-vi.ebv for sequences of Epstein-Barr virus proteins.) All three peptides appear to bind at least three times more antibody from the anti-Sm precipitin positive lupus patient sera than the controls. The antibody binding to these peptides in the lupus patient sera was over half of the antibody binding level found for PPPGMRPP. (SEQ ID NO:4)

Serum samples from lupus patients stored early in the course of the disease process bind only PPPGMRPP (and neighboring peptides) of the 233 possible octapeptides of B/B'.

The analysis of stored sera revealed that a serum sample from an lupus patient stored early in the course of her disease process bound only PPPGMRPP (and neighboring peptides) of the 233 possible octapeptides of B/B' (Figure 4) (James, J.A. et al. J. Exp. Med. 181:453-461 (December 1995)). This repeated PPPGMRPP motif is an early target in three additional patients tested from whom sera were available from early in their disease. In all of these patients PPPGMRPP is the first autoimmune epitope of the Sm B/B' autoantigen against which one can detect antibody binding. One interpretation of this finding is that PPPGMRPP is the first structure bound in at least some of those lupus patients who have anti-Sm autoantibodies. Most, if not nearly all, sera with anti-Sm have more complex binding when they present to their doctors with the illness. At this point no anti-Sm precipitin positive lupus patient who does not have antibodies to PPPGMRPP is known.

Antibodies directed against PPPGMRPP are a significant portion of the anti-Sm response in some patient sera. Five patient sera have been absorbed over columns composed of PPPGMRPP. Four of these sera had anti-Sm and anti-nRNP precipitins and one had an anti-nRNP precipitin without anti-Sm; all five had anti-Ro autoantibodies which were above the normal range. The PPPGMRPP absorption removed 13 to 39% of the patient anti-Sm/nRNP response. These same column absorptions removed less than 10% of the anti-Ro reactivity and over 95% of the anti-PPPGMRPP response. (The proportion of the anti-Sm/nRNP activity absorbed varies with the number of the other octapeptides from Sm B/B' and D proteins bound by immune serum being tested.)

84  
Rabbits immunized with PPPGMRPP developed antibody beyond the peptide of immunization which bound to many other octapeptides in the spliceosome, antinuclear autoantibodies, anti-double stranded DNA autoantibodies and clinical features that suggests the illness known in man as systemic lupus erythematosus (SEQ ID NO:4)

5  
85  
Rabbits were immunized with PPPGMRPP on a MAP<sup>TM</sup> backbone (referred to as ~~PPGMRPP-MAP~~ where the trademark refers only to the MAP<sup>TM</sup>). These animals developed antibody beyond the peptide of immunization which bound to many other octapeptides in the spliceosome. These rabbits usually developed anti-PPPGMRPP antibodies along with anti-Sm and anti-nRNP autoantibodies, antinuclear autoantibodies, anti-double stranded DNA autoantibodies and clinical features that suggests the illness known in man as systemic lupus erythematosus. These rabbits variably had seizures, thrombocytopenia, proteinuria, renal insufficiency, cellular casts in the urine, hypoalbuminemia and alopecia (James, J.A. et al. J. Exp. Med. 181:453-461 (1995)). This has led to a new model of autoimmunity and suggests that a vigorous immune response against this one sequence, PPPGMRPP, is sufficient to induce a specific autoimmune disease in some strains of animals. (SEQ ID NO:4)

20  
25  
30  
86  
PPPGRRP from the Epstein-Barr virus Nuclear Antigen-1 was found when the sequence was searched for sequences similar to PPPGMRPP. To preliminarily test whether it was possible for this sequence to induce spliceosomal autoimmunity, two rabbits were immunized with PPPGRRP-MAP<sup>TM</sup> following the protocol previously used with ~~PPGMRPP-MAP~~. One of the two rabbits developed not only anti-PPPGMRPP antibodies, but also anti-spliceosomal autoantibodies and had B cell epitope spreading to regions of B/B' and D, as evidenced by antibody binding to other octapeptides (Figures 5 and 6). (SEQ ID NO:4)

35  
If a structure from a virus could induce anti-spliceosomal autoantibodies when immunized in an animal then, this is evidence that this structure and the virus is important in the induction of the autoimmune disease associated with anti-spliceosomal autoantibodies, systemic lupus erythematosus.

These data suggest a sequence of events, as follows: (SEQ ID NO: 1)  
infection with Epstein-Barr virus, development of anti-PPPGRRP<sup>1</sup>  
antibodies, development of surface immunoglobulin (and secreted  
antibody) binding both PPPGRRP<sup>1</sup> and PPPGMRPP<sup>1</sup> (the first evidence  
of autoimmunity), antigen processing of the spliceosome bound by  
surface immunoglobulin (or soluble antibody) and presentation of  
the spliceosomal peptides, B cell epitope spreading (and perhaps  
T cell epitope spreading) and clinical features of lupus.  
Epstein-Barr virus infected cells are believed to encourage this  
sequence of events at a number of steps. By substituting another  
antigen for PPPGRRP<sup>1</sup> and another autoantigen for PPPGMRPP<sup>1</sup>, this  
mechanism may generate lupus or another autoimmune disorder by  
following the same general sequence of events.

*Genetic Linkages in Mice to Development of Lupus-like  
Autoimmunity After Immunization with PPPGMRPP-MAP<sup>TM</sup>*  
Experiments in mice have shown that there are some strains  
which develop lupus-like autoimmunity after immunization with  
PPPGMRPP-MAP<sup>TM</sup> and some that do not (J.A. James et al. Arthritis  
Rheum. 38:S226 (1995)). The difference between one responder  
(develops lupus-like autoimmunity) and one nonresponder mouse  
strain has been preliminarily localized to a region on mouse  
chromosome 4 (James, J.A. and Harley, J.B. Arthritis Rheum.  
39:S216 (1996)).

This preliminary finding of linkage in this murine model of  
lupus suggest that the difference between the two parent strains  
of a recombinant inbred set of mice which determines the  
observable B cell epitope spreading and autoimmunity may be found  
at a single gene locus. This region of murine chromosome 4 does  
not contain immunoglobulin genes, T cell receptor genes, nor the  
histocompatibility genes which are highly variable polymorphic  
genes which determine so many other features of the immune  
response. The gene for CD72 is found here which may be  
important. Once the gene responsible for the observed effect is  
identified, then one skilled in the art can provide compositions  
for diagnosis, prevention and treatment of autoimmune disease.  
Beyond this specific example, it is expected that other genes and



gene products will be shown to be important and analogous compositions will be obvious to one skilled in the art.

The host experiences changes after Epstein-Barr virus infection beyond the generation of an immune response. Gene expression and molecular machinery in infected cells is altered by the virus. These changes can be detected by those skilled in the art and used as diagnostics, as well as in the screening and development of therapeutics which are particularly important for individuals affected with or at risk of becoming affected with autoimmune disease.

**Example 2. Association of seroconversion against Epstein-Barr virus and an autoimmune disease.**

Sera from lupus patients compared with controls were first tested for seroconversion against Epstein-Barr virus. There were a number of important considerations in designing these experiments. First, the older and more traditional assays for Epstein-Barr seroconversion were insufficiently specific. These assays are especially unreliable when applied to sera from lupus patients. Antinuclear autoantibodies in lupus sera often interfere with the test for Epstein-Barr seroconversion when an Epstein-Barr virus-infected cell line is used for the test. Both a positive test result for Epstein-Barr virus and for antinuclear antibodies involving the nuclear fluorescence of the cell line. Also, the infected cell line test for Epstein-Barr virus is often not positive in people known to have been infected with Epstein-Barr virus. Others have used such assays in patients with systemic lupus erythematosus with disappointing results (Tsai, Y. et al. Int. Arch. Allergy Immunol. 106:235-240 (1995)).

Second, in the United States as many as 95% of normal adults are infected with and have evidence of seroconversion against Epstein-Barr virus (Evans, A.S. and Niederman, J.C.: Epstein-Barr virus. In Viral Infections in Humans, 3rd ed. Evans, A.S. ed. pp 265-292 (Plenum, New York City 1989)). To detect a difference in adults above the 95% expected baseline of Epstein-Barr virus infection with confidence would require an unreasonably large number of patients and controls. On the other hand, in the

United States, most people contract Epstein-Barr virus before the age of 20 years. Epstein-Barr virus causes infectious mononucleosis, also known as the "kissing disease", in some people. Controls who are children and adolescents will have a lower rate of infection than individuals who are older. By using younger patients and controls, it was thought that one might be able to collect a group of controls where the rate of Epstein-Barr virus seroconversion is less, perhaps less than 75%, thereby a sufficient increase in statistical power to enable this question to be addressed with the resources available.

The use of children and adolescents has nothing to do with anything unique or different about lupus in childhood or adolescence compared with the disease in adults. Rather, the younger cases represent the population in which the hypothesis of a relationship between Epstein-Barr virus infection and autoimmune disease can be most efficiently tested.

The most reliable test for seroconversion is against the Viral Capsid Antigen of Epstein-Barr virus, also called gp340/220 (and abbreviated as EBV-VCA). Solid phase assays are available for antibodies against this surface protein. The purification methods may include other proteins and antigens from Epstein-Barr virus, but as long as a substantial proportion of the preparation is composed of this surface antigen of Epstein-Barr virus, then the assay should be sensitive for an immune response directed against Epstein-Barr virus. The procedure used to isolate the antigen used in the commercial ELISA (enzyme-linked immunosorbent assay) kits has been described in detail (Qualtiere, L.F. and Pearson, G.R. Virology 102:360-369 (1980)). Antibodies binding to the Epstein-Barr virus surface protein are thought to occur in virtually everyone who is infected with Epstein-Barr virus. An assay for antibodies binding to the Epstein-Barr virus Viral Capsid Antigen (EBV-VCA) is marketed by only two manufacturers in the United States. The assay first used was manufactured by Clark Laboratories, Inc. (Jamestown, New York). This firm was recently acquired by Wampole Laboratories, a division of Carter-Wallace, Inc. (Cranbury, New Jersey) who has continued to provide

the same viral assays. The manufacturer's instructions were followed in measuring the anti-VCA Epstein-Barr virus IgG antibodies in lupus patients and their controls, all of whom tested are under 20 years old.

5 The results are striking. Of the 134 tested sera from controls, 96 have IgG antibodies that bind at a sufficient level to conclude that these individuals have seroconverted. The sera from the lupus patients in this experiment are nearly all seropositive at 102 of 103 or over 99% (Table 1).

08791295-011397

Table 1. Seroconversion frequencies in pediatric lupus and controls for IgG binding to Epstein Barr virus viral capsid antigen (EBV-VCA).

Anti-Epstein-Barr virus Viral Capsid Antigen IgG  
(# seroconverted (total tested))

Oklahoma cases and controls

Lupus cases	45 (45)
Controls	53 (76)
Odds ratio	38
$\chi^2$	15.1
p	<0.0001

San Diego and New York City cases and controls

Lupus cases	57 (58)
Controls	43 (58)
Odds ratio	20
$\chi^2$	14.1
p	<0.001

Combined data from lupus cases and controls

Lupus cases	102 (103)
Controls	96 (134)
Odds ratio	40.4
$\chi^2$	31.6
p	<0.00000001

Odds ratios were calculated from the contingency tables. A  $\chi^2$  test was used to assess significance. The assays for all viral antigens were obtained from Clark Laboratories (Jamestown, NY) now a division of Wampole Laboratories (Cranberry, New Jersey) and were performed according to the instructions provided by the manufacturer. The surface antigen is partially purified for the anti-Epstein-Barr virus Viral Capsid Antigen antibody assay (Qualtiere, L.F. and Pearson, G.R. Virology 102:360-369 (1980)).

The quantitative level of binding to the Epstein-Barr virus Viral Capsid Antigen (as measured relative to the standardized

calibrator and as recommended by the manufacturer) also show substantial quantitative differences between lupus cases and controls (Table 2).

**Table 2. Binding of IgG in sera from pediatric lupus patients and their controls to the Viral Capsid Antigen from Epstein Barr virus (EBV).**

<b>Epstein-Barr virus Viral Capsid Antigen (International Standardized Ratios (ISRs))</b>	
Lupus cases (n=103) (ISRs)	4.368
Controls (n=134) (ISRs)	2.087
t-test	14.82
probability (p value)	<<0.001

Assay is performed as presented in Table 1. Data are presented as International Standardized Ratios (ISRs) which are calculated as described by the manufacturer. The average binding of each tested serum is divided by the product of the calibrator and a lot-specific factor. This allows for standardization across assays. Values <0.9 are considered negative and values >1.1 are considered positive. Values between 0.9 and 1.1 are equivocal and are either retested or are assigned to the not positive or negative group. (No patient serum tested to date has had a result in this equivocal range.)

If the suspected relationship between systemic lupus erythematosus held only for the lupus patients who were anti-Sm positive, then the proportion of lupus patients who had developed the anti-Epstein-Barr virus surface molecule (gp340/220) antibody would be increased over controls only modestly, since the proportion of lupus patients with anti-Sm in their sera rarely exceeds 30%. The remaining lupus patients would have been expected to have the normal control frequency of Epstein-Barr virus seroconversion.

Patients with lupus have such different clinical manifestations from patient to patient, that many investigators suspect that there are many different causes of this disease. Any given particular etiology causing lupus would then be expected to account for only a proportion of the cases. For

example, assume that lupus, as observed in the clinics, happens to have two different origins, one from a virus such as Epstein-Barr virus and another from the intrinsic genetic program of some patients. Further assume that the Epstein-Barr virus only  
5 accounts for the individual patients who have anti-Sm or anti-nRNP autoantibodies and that this is half of the lupus patients (which is approximately true). Under this scenario, one would expect to see a statistical effect of increased Epstein-Barr virus infection in the group of lupus patients, as a whole, when  
10 compared to controls, but the effect would be driven by only half of the patients. Consequently, the proportion of lupus patients with evidence of Epstein-Barr virus infection would be raised, from 71% to 85.5%, but not to the 99% level that has been observed. This 99% rate is the level that would be expected if over 95% of the lupus cases in the collection have been caused by Epstein-barr virus.

461226 - 011307  
20  
The association is consistent with Epstein-Barr virus infection being a necessary condition before this autoimmune disease can develop in nearly all lupus patients. Assuming Epstein-Barr virus accounts for nearly all cases of lupus, as suggested by the data, then this virus is capable of mediating the various autoimmune manifestations found in lupus beyond just the relationship with anti-Sm antigen originally postulated. Autoantigens in lupus patients are generally thought to be found  
25 throughout the tissues of the host and not to be restricted to a single tissue. This is true for Sm, nRNP, Ro, La, P, DNA, and RNA, among many other autoantigens in lupus. Some patients may have antibodies binding other autoantigens which are tissue specific, such as thyroglobulin, but these seem to be less  
30 important in this disease. Virtually all active lupus patients have a positive antinuclear antibody, suggesting that at least one of the autoantigens bound by patient autoantibody is directed against a cellular constituent shared between tissues.

This mechanism already allows for a large number of  
35 autoantigens. If all of these autoimmune manifestations are related to Epstein-Barr virus infection, then it is probable that

other autoimmune disorders are related also. Only a change in the location of the antigen in the host would be required to generate another disorder, among other mechanisms, for Epstein-Barr virus to be a necessary, but not sufficient condition for the development of other autoimmune diseases. For example, a similar mechanism could operate for thyroglobulin or thyroid peroxidase, leading to a tissue specific autoimmune disorder, autoimmune thyroid disease. Again Epstein-Barr virus infection would be a requirement before most patients could develop the disease.

Interestingly, autoimmune thyroid <sup>disease</sup>~~disease~~ and lupus sometimes coexist in the same patient. There are those who have suspected that there is a relationship between systemic lupus erythematosus and autoimmune thyroid disease (R. H. Scofield Clin. Exp. Rheum. 14:321-330 (1996)). Under these circumstances, Epstein-Barr virus could be fundamental to both disorders. Other features of the environment, virus strain differences, genetics of the host, hormone and cytokine status, recent and remote immune history, and the nature and course of infection of the Epstein-Barr virus in the host, among other variables, probably then determine how the autoimmunity, as a consequence of Epstein-Barr virus infection, finds expression.

**Example 3. Experiments to evaluate trivial and artifactual explanations for the association of Epstein-Barr virus serologic positivity with an autoimmune disease related to the source of the test for anti-Epstein-Barr Viral Capsid Antigen IgG antibodies.**

Perhaps the association observed is explained by some artefact of the manufacturing process. Since there are two manufacturers who separately prepare the Viral Capsid Antigen and who use somewhat different procedures, a subset of the lupus patients and controls were evaluated using the test manufactured by Gull Laboratories (Salt Lake City, Utah).

The results from the Gull test for Epstein-Barr seropositivity were virtually identical to those obtained with

tests manufactured by Clark Laboratories, Inc. Of the 43 patient sera tested, all were positive for Epstein-Barr virus IgG antibodies who were positive by the Clark assay. The one patient without antibodies to Epstein-Barr virus Viral Capsid Antigen by the Clark assay also had no detectable antibodies to Epstein-Barr virus Viral Capsid Antigen by the Gull assay. Of the 47 control sera tested, none were negative that were previously positive and four were positive in the Gull anti-Epstein-Barr virus Viral Capsid Antigen assay that were previously negative.

This experiment established that the results in Example 1 were not explained by lot variation or an unusual property of the preparation used to detect anti-Epstein-Barr virus surface antigen.

**Example 4. Experiments to evaluate trivial and artifactual explanations for the association of Epstein-Barr virus serologic positivity with an autoimmune disease related to cross-reactivity of autoantibodies with anti-Epstein-Barr surface antigen IgG antibodies.**

Next, five of the lupus patients with high levels of anti-Sm autoantibodies were arbitrarily selected. The anti-spliceosomal autoantibodies were absorbed from their sera and their antinuclear antibody titer reevaluated by immunofluorescence, and anti-Sm/nRNP and anti-Ro antibodies by solid phase assay (following previous methods (Gaither, K.K. et al. J. Clin. Invest. 79:841-846 (1987); Harley, J.B. et al. Arthritis Rheum. 29:196-206 (1986)), and anti-Epstein-Barr virus Viral Capsid Antigen IgG antibodies.

The spliceosomal antigen preparation is affinity purified from bovine tissue and contains both the Sm and nRNP specificities and, hence, is referred to as Sm/nRNP. This is the same preparation discussed herein for solid phase assays against Sm/nRNP. The previous method for the solid phase anti-Sm/nRNP assay (Gaither, K.K. et al. J. Clin. Invest. 79:841-846 (1987); Harley, J.B. et al. Arthritis Rheum. 29:196-206 (1986)) was



altered by omitting inhibition with purified Sm/nRNP antigen.  
The same alteration was made in the anti-Ro assay.

These data show that the anti-Sm was reduced by at least 90%  
in each serum, as expected. In addition, the antinuclear  
antibodies were reduced by an average of 97%. Meanwhile, neither  
the anti-Ro nor the anti-Epstein-Barr virus surface antigen IgG  
antibodies were substantially reduced. The anti-Epstein-Barr  
virus surface antigen IgG antibodies were reduced by an average  
of 8%.

These data show that neither anti-Ro nor anti-Epstein-Barr  
virus Viral Capsid Antigen IgG cross-reacts to any significant  
extent with the anti-Sm. If they had, then these antibodies  
would also have been removed by the absorption. In addition, the  
reduction of the antinuclear antibodies in the absorbed sera by  
97% suggests that the anti-Sm/nRNP is the major autoantigen in  
these lupus patients.

**Example 5. Experiments to evaluate trivial and artifactual  
explanations for the association of Epstein-Barr  
virus serologic positivity with an autoimmune  
disease related a general polyclonality of  
autoantibodies in lupus sera.**

It was possible that lupus sera may nonspecifically bind to  
any viral antigen. To test this possibility, the same patient  
and control sera were evaluated for antibodies against four other  
Herpes viruses: Cytomegalovirus; Herpes simplex, type 1; Herpes  
simplex, Type 2; and Varicella zoster (Table 3).

Table 3. Seroconversion frequencies in pediatric lupus and controls for IgG binding to cytomegalovirus antigen (CMV), Herpes simplex type 1 antigen (HSV-1), Herpes simplex virus type 2 antigen (HSV-2), and varicella zoster virus (VZV) antigens.

	CMV	HSV-1	HSV-2	VZV
<u>Oklahoma</u>				
Lupus cases	20 (45)	30 (45)	21 (45)	42 (45)
Controls	21 (76)	32 (76)	22 (76)	71 (76)
Odds ratio	2.1	2.75	2.14	0.99
$\chi^2$	3.57	6.82	3.84	0.0
p	>0.05	<0.01	0.05	0.99

San Diego and New York City

Lupus cases	18 (58)	33 (58)	32 (58)	46 (58)
Controls	12 (57)	31 (57)	23 (57)	45 (57)
Odds ratio	1.69	1.11	1.82	1.02
$\chi^2$	1.49	0.73	2.53	0.0
p	0.22	0.79	0.11	0.96

The odds ratios were calculated from the contingency tables. A  $\chi^2$  test was used to assess significance. The assays for all viral antigens were obtained from Clark Laboratories (Jamestown, NY) now a division of Wampole Laboratories (Cranberry, New Jersey) and were performed according to the instructions provided by the manufacturer. Whole inactivated virus is used for the CMV, HSV-1, HSV-2 and VZV antibody assays.

There is no consistent difference (at the level of  $p < 0.05$ ) between the controls and the lupus patients for the frequency of seroconversion against any of the viruses tested except for Epstein-Barr virus. There is a small and significant difference between the cases and controls for IgG autoantibodies against herpes simplex, Types 1 and 2 in the Oklahoma group (Table 3). Its inconsistency is typical of the results of other assays for seroconversion rates in lupus. Of these, only the difference between cases and controls for antibodies against Herpes Simplex virus Type 2 is significant across the entire collection, odds ratio is 2.07 ( $\chi^2 = 7.39$ ,  $p = 0.006$ ). The low, but significant, odds

ratio, suggests that the contribution of this virus is likely to be small. Some of the social behaviors that increase the risk of infection with Epstein-Barr virus probably secondarily also increase the risk of contracting Herpes Simplex virus, Type 2.

5 The modestly, and usually insignificantly, increased odds ratios for the seroprevalence of cytomegalovirus and Herpes simplex Types 1 and 2 in lupus patients compared to controls may reflect the increased levels of binding of antibodies found  
D 10 ~~in other~~ studies (Hollinger, F.B. et al. Bact. Proc. 131:174 (1970); Phillips, P.E. and Christian, C.L. Science 168:982-4 (1970); Hurd, E.R. et al. Arthritis Rheum. 13:724-33 (1970)).

In conclusion, these control assays do not support the position that the difference in the seroconversion rate between lupus patients and their collected controls is the result of a nonspecific binding to viral antigens.

25  
30  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995  
1000  
1005  
1010  
1015  
1020  
1025  
1030  
1035  
1040  
1045  
1050  
1055  
1060  
1065  
1070  
1075  
1080  
1085  
1090  
1095  
1100  
1105  
1110  
1115  
1120  
1125  
1130  
1135  
1140  
1145  
1150  
1155  
1160  
1165  
1170  
1175  
1180  
1185  
1190  
1195  
1200  
1205  
1210  
1215  
1220  
1225  
1230  
1235  
1240  
1245  
1250  
1255  
1260  
1265  
1270  
1275  
1280  
1285  
1290  
1295  
1300  
1305  
1310  
1315  
1320  
1325  
1330  
1335  
1340  
1345  
1350  
1355  
1360  
1365  
1370  
1375  
1380  
1385  
1390  
1395  
1400  
1405  
1410  
1415  
1420  
1425  
1430  
1435  
1440  
1445  
1450  
1455  
1460  
1465  
1470  
1475  
1480  
1485  
1490  
1495  
1500  
1505  
1510  
1515  
1520  
1525  
1530  
1535  
1540  
1545  
1550  
1555  
1560  
1565  
1570  
1575  
1580  
1585  
1590  
1595  
1600  
1605  
1610  
1615  
1620  
1625  
1630  
1635  
1640  
1645  
1650  
1655  
1660  
1665  
1670  
1675  
1680  
1685  
1690  
1695  
1700  
1705  
1710  
1715  
1720  
1725  
1730  
1735  
1740  
1745  
1750  
1755  
1760  
1765  
1770  
1775  
1780  
1785  
1790  
1795  
1800  
1805  
1810  
1815  
1820  
1825  
1830  
1835  
1840  
1845  
1850  
1855  
1860  
1865  
1870  
1875  
1880  
1885  
1890  
1895  
1900  
1905  
1910  
1915  
1920  
1925  
1930  
1935  
1940  
1945  
1950  
1955  
1960  
1965  
1970  
1975  
1980  
1985  
1990  
1995  
2000  
2005  
2010  
2015  
2020  
2025  
2030  
2035  
2040  
2045  
2050  
2055  
2060  
2065  
2070  
2075  
2080  
2085  
2090  
2095  
2100  
2105  
2110  
2115  
2120  
2125  
2130  
2135  
2140  
2145  
2150  
2155  
2160  
2165  
2170  
2175  
2180  
2185  
2190  
2195  
2200  
2205  
2210  
2215  
2220  
2225  
2230  
2235  
2240  
2245  
2250  
2255  
2260  
2265  
2270  
2275  
2280  
2285  
2290  
2295  
2300  
2305  
2310  
2315  
2320  
2325  
2330  
2335  
2340  
2345  
2350  
2355  
2360  
2365  
2370  
2375  
2380  
2385  
2390  
2395  
2400  
2405  
2410  
2415  
2420  
2425  
2430  
2435  
2440  
2445  
2450  
2455  
2460  
2465  
2470  
2475  
2480  
2485  
2490  
2495  
2500  
2505  
2510  
2515  
2520  
2525  
2530  
2535  
2540  
2545  
2550  
2555  
2560  
2565  
2570  
2575  
2580  
2585  
2590  
2595  
2600  
2605  
2610  
2615  
2620  
2625  
2630  
2635  
2640  
2645  
2650  
2655  
2660  
2665  
2670  
2675  
2680  
2685  
2690  
2695  
2700  
2705  
2710  
2715  
2720  
2725  
2730  
2735  
2740  
2745  
2750  
2755  
2760  
2765  
2770  
2775  
2780  
2785  
2790  
2795  
2800  
2805  
2810  
2815  
2820  
2825  
2830  
2835  
2840  
2845  
2850  
2855  
2860  
2865  
2870  
2875  
2880  
2885  
2890  
2895  
2900  
2905  
2910  
2915  
2920  
2925  
2930  
2935  
2940  
2945  
2950  
2955  
2960  
2965  
2970  
2975  
2980  
2985  
2990  
2995  
3000  
3005  
3010  
3015  
3020  
3025  
3030  
3035  
3040  
3045  
3050  
3055  
3060  
3065  
3070  
3075  
3080  
3085  
3090  
3095  
3100  
3105  
3110  
3115  
3120  
3125  
3130  
3135  
3140  
3145  
3150  
3155  
3160  
3165  
3170  
3175  
3180  
3185  
3190  
3195  
3200  
3205  
3210  
3215  
3220  
3225  
3230  
3235  
3240  
3245  
3250  
3255  
3260  
3265  
3270  
3275  
3280  
3285  
3290  
3295  
3300  
3305  
3310  
3315  
3320  
3325  
3330  
3335  
3340  
3345  
3350  
3355  
3360  
3365  
3370  
3375  
3380  
3385  
3390  
3395  
3400  
3405  
3410  
3415  
3420  
3425  
3430  
3435  
3440  
3445  
3450  
3455  
3460  
3465  
3470  
3475  
3480  
3485  
3490  
3495  
3500  
3505  
3510  
3515  
3520  
3525  
3530  
3535  
3540  
3545  
3550  
3555  
3560  
3565  
3570  
3575  
3580  
3585  
3590  
3595  
3600  
3605  
3610  
3615  
3620  
3625  
3630  
3635  
3640  
3645  
3650  
3655  
3660  
3665  
3670  
3675  
3680  
3685  
3690  
3695  
3700  
3705  
3710  
3715  
3720  
3725  
3730  
3735  
3740  
3745  
3750  
3755  
3760  
3765  
3770  
3775  
3780  
3785  
3790  
3795  
3800  
3805  
3810  
3815  
3820  
3825  
3830  
3835  
3840  
3845  
3850  
3855  
3860  
3865  
3870  
3875  
3880  
3885  
3890  
3895  
3900  
3905  
3910  
3915  
3920  
3925  
3930  
3935  
3940  
3945  
3950  
3955  
3960  
3965  
3970  
3975  
3980  
3985  
3990  
3995  
4000  
4005  
4010  
4015  
4020  
4025  
4030  
4035  
4040  
4045  
4050  
4055  
4060  
4065  
4070  
4075  
4080  
4085  
4090  
4095  
4100  
4105  
4110  
4115  
4120  
4125  
4130  
4135  
4140  
4145  
4150  
4155  
4160  
4165  
4170  
4175  
4180  
4185  
4190  
4195  
4200  
4205  
4210  
4215  
4220  
4225  
4230  
4235  
4240  
4245  
4250  
4255  
4260  
4265  
4270  
4275  
4280  
4285  
4290  
4295  
4300  
4305  
4310  
4315  
4320  
4325  
4330  
4335  
4340  
4345  
4350  
4355  
4360  
4365  
4370  
4375  
4380  
4385  
4390  
4395  
4400  
4405  
4410  
4415  
4420  
4425  
4430  
4435  
4440  
4445  
4450  
4455  
4460  
4465  
4470  
4475  
4480  
4485  
4490  
4495  
4500  
4505  
4510  
4515  
4520  
4525  
4530  
4535  
4540  
4545  
4550  
4555  
4560  
4565  
4570  
4575  
4580  
4585  
4590  
4595  
4600  
4605  
4610  
4615  
4620  
4625  
4630  
4635  
4640  
4645  
4650  
4655  
4660  
4665  
4670  
4675  
4680  
4685  
4690  
4695  
4700  
4705  
4710  
4715  
4720  
4725  
4730  
4735  
4740  
4745  
4750  
4755  
4760  
4765  
4770  
4775  
4780  
4785  
4790  
4795  
4800  
4805  
4810  
4815  
4820  
4825  
4830  
4835  
4840  
4845  
4850  
4855  
4860  
4865  
4870  
4875  
4880  
4885  
4890  
4895  
4900  
4905  
4910  
4915  
4920  
4925  
4930  
4935  
4940  
4945  
4950  
4955  
4960  
4965  
4970  
4975  
4980  
4985  
4990  
4995  
5000  
5005  
5010  
5015  
5020  
5025  
5030  
5035  
5040  
5045  
5050  
5055  
5060  
5065  
5070  
5075  
5080  
5085  
5090  
5095  
5100  
5105  
5110  
5115  
5120  
5125  
5130  
5135  
5140  
5145  
5150  
5155  
5160  
5165  
5170  
5175  
5180  
5185  
5190  
5195  
5200  
5205  
5210  
5215  
5220  
5225  
5230  
5235  
5240  
5245  
5250  
5255  
5260  
5265  
5270  
5275  
5280  
5285  
5290  
5295  
5300  
5305  
5310  
5315  
5320  
5325  
5330  
5335  
5340  
5345  
5350  
5355  
5360  
5365  
5370  
5375  
5380  
5385  
5390  
5395  
5400  
5405  
5410  
5415  
5420  
5425  
5430  
5435  
5440  
5445  
5450  
5455  
5460  
5465  
5470  
5475  
5480  
5485  
5490  
5495  
5500  
5505  
5510  
5515  
5520  
5525  
5530  
5535  
5540  
5545  
5550  
5555  
5560  
5565  
5570  
5575  
5580  
5585  
5590  
5595  
5600  
5605  
5610  
5615  
5620  
5625  
5630  
5635  
5640  
5645  
5650  
5655  
5660  
5665  
5670  
5675  
5680  
5685  
5690  
5695  
5700  
5705  
5710  
5715  
5720  
5725  
5730  
5735  
5740  
5745  
5750  
5755  
5760  
5765  
5770  
5775  
5780  
5785  
5790  
5795  
5800  
5805  
5810  
5815  
5820  
5825  
5830  
5835  
5840  
5845  
5850  
5855  
5860  
5865  
5870  
5875  
5880  
5885  
5890  
5895  
5900  
5905  
5910  
5915  
5920  
5925  
5930  
5935  
5940  
5945  
5950  
5955  
5960  
5965  
5970  
5975  
5980  
5985  
5990  
5995  
6000  
6005  
6010  
6015  
6020  
6025  
6030  
6035  
6040  
6045  
6050  
6055  
6060  
6065  
6070  
6075  
6080  
6085  
6090  
6095  
6100  
6105  
6110  
6115  
6120  
6125  
6130  
6135  
6140  
6145  
6150  
6155  
6160  
6165  
6170  
6175  
6180  
6185  
6190  
6195  
6200  
6205  
6210  
6215  
6220  
6225  
6230  
6235  
6240  
6245  
6250  
6255  
6260  
6265  
6270  
6275  
6280  
6285  
6290  
6295  
6300  
6305  
6310  
6315  
6320  
6325  
6330  
6335  
6340  
6345  
6350  
6355  
6360  
6365  
6370  
6375  
6380  
6385  
6390  
6395  
6400  
6405  
6410  
6415  
6420  
6425  
6430  
6435  
6440  
6445  
6450  
6455  
6460  
6465  
6470  
6475  
6480  
6485  
6490  
6495  
6500  
6505  
6510  
6515  
6520  
6525  
6530  
6535  
6540  
6545  
6550  
6555  
6560  
6565  
6570  
6575  
6580  
6585  
6590  
6595  
6600  
6605  
6610  
6615  
6620  
6625  
6630  
6635  
6640  
6645  
6650  
6655  
6660  
6665  
6670  
6675  
6680  
6685  
6690  
6695  
6700  
6705  
6710  
6715  
6720  
6725  
6730  
6735  
6740  
6745  
6750  
6755  
6760  
6765  
6770  
6775  
6780  
6785  
6790  
6795  
6800  
6805  
6810  
6815  
6820  
6825  
6830  
6835  
6840  
6845  
6850  
6855  
6860  
6865  
6870  
6875  
6880  
6885  
6890  
6895  
6900  
6905  
6910  
6915  
6920  
6925  
6930  
6935  
6940  
6945  
6950  
6955  
6960  
6965  
6970  
6975  
6980  
6985  
6990  
6995  
7000  
7005  
7010  
7015  
7020  
7025  
7030  
7035  
7040  
7045  
7050  
7055  
7060  
7065  
7070  
7075  
7080  
7085  
7090  
7095  
7100  
7105  
7110  
7115  
7120  
7125  
7130  
7135  
7140  
7145  
7150  
7155  
7160  
7165  
7170  
7175  
7180  
7185  
7190  
7195  
7200  
7205  
7210  
7215  
7220  
7225  
7230  
7235  
7240  
7245  
7250  
7255  
7260  
7265  
7270  
7275  
7280  
7285  
7290  
7295  
7300  
7305  
7310  
7315  
7320  
7325  
7330  
7335  
7340  
7345  
7350  
7355  
7360  
7365  
7370  
7375  
7380  
7385  
7390  
7395  
7400  
7405  
7410  
7415  
7420  
7425  
7430  
7435  
7440  
7445  
7450  
7455  
7460  
7465  
7470  
7475  
7480  
7485  
7490  
7495  
7500  
7505  
7510  
7515  
7520  
7525  
7530  
7535  
7540  
7545  
7550  
7555  
7560  
7565  
7570  
7575  
7580  
7585  
7590  
7595  
7600  
7605  
7610  
7615  
7620  
7625  
7630  
7635  
7640  
7645  
7650  
7655  
7660  
7665  
7670  
7675  
7680  
7685  
7690  
7695  
7700  
7705  
7710  
7715  
7720  
7725  
7730  
7735  
7740  
7745  
7750  
7755  
7760  
7765  
7770  
7775  
7780  
7785  
7790  
7795  
7800  
7805  
7810  
7815  
7820  
7825  
7830  
7835  
7840  
7845  
7850  
7855  
7860  
7865  
7870  
7875  
7880  
7885  
7890  
7895  
7900  
7905  
7910  
7915  
7920  
7925  
7930  
7935  
7940  
7945  
7950  
7955  
7960  
7965  
7970  
7975  
7980  
7985  
7990  
7995  
8000  
8005  
8010  
8015  
8020  
8025  
8030  
8035  
8040  
8045  
8050  
8055  
8060  
8065  
8070  
8075  
8080  
8085  
8090  
8095  
8100  
8105  
8110  
8115  
8120  
8125  
8130  
8135  
8140  
8145  
8150  
8155  
8160  
8165  
8170  
8175  
8180  
8185  
8190  
8195  
8200  
8205  
8210  
8215  
8220  
8225  
8230  
8235  
8240  
8245  
8250  
8255  
8260  
8265  
8270  
8275  
8280  
8285  
8290  
8295  
8300  
8305  
8310  
8315  
8320  
8325  
8330  
8335  
8340  
8345  
8350  
8355  
8360  
8365  
8370  
8375  
8380  
8385  
8390  
8395  
8400  
8405  
8410  
8415  
8420  
8425  
8430  
8435  
8440  
8445  
8450  
8455  
8460  
8465  
8470  
8475  
8480  
8485  
8490  
8495  
8500  
8505  
8510  
8515  
8520  
8525  
8530  
8535  
8540  
8545  
8550  
8555  
8560  
8565  
8570  
8575  
8580  
8585  
8590  
8595  
8600  
8605  
8610  
8615  
8620  
8625  
8630  
8635  
8640  
8645  
8650  
8655  
8660  
8665  
8670  
8675  
8680  
8685  
8690  
8695  
8700  
8705  
8710  
8715  
8720  
8725  
8730  
8735  
8740  
8745  
8750  
8755  
8760  
8765  
8770  
8775  
8780  
8785  
8790  
8795  
8800  
8805  
8810  
8815  
8820  
8825  
8830  
8835  
8840  
8845  
8850  
8855  
8860  
8865  
8870  
8875  
8880  
8885  
8890  
8895  
8900  
8905  
8910  
8915  
8920  
8925  
8930  
8935  
8940  
8945  
8950  
8955  
8960  
8965  
8970  
8975  
8980  
8985  
8990  
8995  
9000  
9005  
9010  
9015  
9020  
9025  
9030  
9035  
9040  
9045  
9050  
9055  
9060  
9065  
9070  
9075  
9080  
9085  
9090  
9095  
9100  
9105  
9110  
9115  
9120  
9125  
9130  
9135  
9140  
9145  
9150  
9155  
9160  
9165  
9170  
9175  
9180  
9185  
9190  
9195  
9200  
9205  
9210  
9215  
9220  
9225  
9230  
9235  
9240  
9245  
9250  
9255  
9260  
9265  
9270  
9275  
9280  
9285  
9290  
9295  
9300  
9305  
9310  
9315  
9320  
9325  
9330  
9335  
9340  
9345  
9350  
9355  
9360  
9365  
9370  
9375  
9380  
9385  
9390  
9395  
9400  
9405  
9410  
9415  
9420  
9425  
9430  
9435  
9440  
9445  
9450  
9455  
9460  
9465  
9470  
9475  
9480  
9485  
9490  
9495  
9500  
9505  
9510  
9515  
9520  
9525  
9530  
9535  
9540  
9545  
9550  
9555  
9560  
9565  
9570  
9575  
9580  
9585  
9590  
9595  
9600  
9605  
9610  
9615  
9620  
9625  
9630  
9635  
9640  
9645  
9650  
9655  
9660  
9665  
9670  
9675  
9680  
9685  
9690  
9695  
9700  
9705  
9710  
9715  
9720  
9725  
9730  
9735  
9740  
9745  
9750  
9755  
9760  
9765  
9770  
9775  
9780  
9785  
9790  
9795  
9800  
9805  
9810  
9815  
9820  
9825  
9830  
9835  
9840  
9845  
9850  
9855  
9860  
9865  
9870  
9875  
9880  
9885  
9890  
9895  
9900  
9905  
9910  
9915  
9920  
9925  
9930  
9935  
9940  
9945  
9950  
9955  
9960  
9965  
9970  
9975  
9980  
9985  
9990  
9995  
10000  
10005  
10010  
10015  
10020  
10025  
10030  
10035  
10040  
10045  
10050  
10055  
10060  
10065  
10070  
10075  
10080  
10085  
10090  
10095  
10100  
10105  
10110  
10115  
10120  
10125  
10130  
10135  
10140  
10145  
10150  
10155  
10160  
10165  
10170

serology, a method altered from the DNA based assays already available was developed. The BamHI W repeat nucleic acid sequence of the Epstein-Barr virus genome contains a sequence repeated 11 times that others have used to detect the virus (Saito et al., J. Exp. Med. 169:2191-2198, 1989; and Miyashita et al., Cell 80:593-601, 1995). These assays were evaluated in an effort to increase specificity. Two mcg DNA specimens were isolated from peripheral blood mononuclear cells. Quantitation of DNA was based both upon optical density measurements at 240 and 260 nm and quantitation of ethidium bromide fluorescence in agarose gel relative to known DNA standards. Where available, six reactions, each containing 2 mcg of mononuclear cell DNA, were evaluated from each subject. The primers and the probe used are given in Table 4.

**Table 4. Primers and probe used to expand Epstein-Barr virus DNA from the BamHI W repeat.** (SEQ ID NO: 10)

Forward - 5'-CCAGAGGTAAGTGGACTT-3'

Reverse - 5'-~~CCAGAGGTAAGTGGACTT~~-3'

Probe - 5'-AAGACGATTCGGGTTG (TGAGGTGGTGTGGGTCCGTGTGTGATGTGTGTGGGTGGGCAG)\*-3' (SEQ ID NO: 12)

\* The <sup>32</sup>P-dCTP label is incorporated into the portion of the sequence in parentheses.

The polymerase chain reaction for each DNA specimen was run in a final volume of 75 µl with 50 mM KCl, 10 mM Tris-HCl at pH 8.0, 1.5 mM MgCl<sub>2</sub>, 0.1% Triton X-100, 0.2 mM each dNTP, 0.5 µM primer, and 2.5 U Taq DNA polymerase. A hot start protocol was followed using Ampliwax PCR Gems™ (Perkin-Elmer, Branchburg, N.J.). The thermocycler was programmed for the following temperature cycles: (95°C for 2 min, 57°C for 1 min, 72°C for 1 min) twice, (94°C for 1 min, 55°C for 1 min, 72°C for 45 sec) 31 times, 72°C for 5 min.

The primers variably expanded four bands from found at approximately 92, 122, ~500, and ~700 base pairs. All four were cloned into the pCRII vector (Invitrogen, San Diego, California) and at least partially sequenced. The 92, ~500, and ~700 base

pair products of the polymerase chain reaction are not in the GeneBank database. They are likely to represent human sequence from a region of the genome that has not yet been sequenced.

The 122 base pair product was similarly cloned and sequenced. The sequence obtained exactly matched the sequence predicted from the Epstein-Barr virus *Bam*HI W repeat, which is repeated 11 times and found at positions 14614-14735, 17686-17807, 20758-20879, 23830-23951, 26902-27023, 29974-30095, 33046-33167, 36118-36239, 39190-39311, 42262-42383, and 45334-45455 of the Epstein-Barr DNA sequence (GenBank accession number: v01555). Only this product of the polymerase chain reaction, found at 122 base pairs, hybridized to the radiolabeled probe.

The existing assays for Epstein-Barr DNA were modified so that the assay applied would be as sensitive and specific as the serologic assays are for infection in normal individuals. To achieve this goal a total of 12 mcg of mononuclear DNA in six polymerase chain reactions were evaluated. A long probe was used so that very small amounts of expanded *Bam*HI W fragment DNA can be detected.

This assay was used in 50 subjects. Of these, 38 individuals that had positive Epstein-Barr virus Viral Capsid Antigen ("EBV-VCA") IgG antibodies (as determined by the assay from Clark Laboratories, Inc.) also expanded the predicted 122 base pair DNA fragment which hybridized to the *Bam*HI W probe from the Epstein-Barr viral DNA sequence. Epstein-Barr virus DNA was recovered in three other subjects whose serologic assays were not positive. In one of these subjects the serologic results for Epstein-Barr virus Viral Capsid Antigen IgG was close to the positive range, as determined in comparison to the calibrator sera provided by the manufacturer. One control had evidence of antibodies to EBV-VCA but no evidence of EBV-DNA as determined by the method described herein. Neither anti-Epstein-Barr virus Viral Capsid Antigen IgG antibodies nor Epstein-Barr virus DNA was detected in the other seven subjects tested.

The new assay has high sensitivity and specificity relative to the serologic analysis. Actually, the direct detection of

viral DNA, if true, is a more proximal indication of infection than is seroconversion and is probably a better standard for viral infection than is serology.

5 The detection of Epstein-Barr virus DNA has proven to be sufficiently reliable for application to the evaluation of the frequency of Epstein-Barr virus infection in lupus and their controls. A matched case-control design was followed. Cases of lupus in children and adolescents (<20 years old) had been enrolled into this experiment as they were encountered along with a sex, age (within two years), and race matched control. To control for socioeconomic factors and likelihood of viral exposure, each case was asked to provide a relative or friend to serve as a control. Relatives were preferred over friends. The investigators selected controls for eight cases who did not provide their own control.

10 The data were evaluated with the Exact Binomial Test for Matched Data. Of the 25 lupus cases, all 25 had been infected by Epstein-Barr virus as determined by the assay for viral DNA and also had antibodies directed against Epstein-Barr Viral Capsid Antigen. Of the 25 matched controls, 17 had been infected by Epstein-Barr virus as determined by the assay for viral DNA. Of the 17 with viral DNA, 14 had IgG antibodies directed against Epstein-Barr virus Viral Capsid Antigen and three did not. One control had no detectable DNA from the virus by the assay and yet had anti-Epstein-Barr virus Viral Capsid Antigen IgG antibodies.

25 To evaluate these data from the most conservative perspective, subjects were required to be negative in both assays in order to conclude that a particular subject had not been infected with Epstein-Barr virus. By these criteria seven of the matched pairs were discordant for infection. All seven of these discordant pairs are discordant in the same way, Epstein-Barr virus infection had occurred in the case and not in the control, which is very unlikely to have occurred by chance ( $p < 0.008$ ) (Table 5).

Table 5. Matched case-control evaluation of the presence of Epstein-Barr DNA in peripheral blood mononuclear cells from lupus patients (cases) and controls matched for sex, race, and age (+/- 2 years).

Epstein-Barr virus infection		
Lupus Cases	Matched Control	
+	+	18
+	-	7
-	+	0
-	-	0

The probability of this distribution occurring by chance is  $p=0.0078$  by the Exact Binomial Test for Matched Data. The odds ratio for this distribution cannot be calculated because of the zero cell. If the usual adjustment of 0.5 is made then the odds ratio is estimated to be  $=\backslash > 13$ .

These results dramatically confirm the serologic data. Collectively, these data, the serologic results and the control experiments establish that there is a virtually complete association of Epstein-Barr virus infection with systemic lupus erythematosus.

**Example 8.** Antibodies against some Epstein-Barr virus structures are associated with the presence of an autoimmune disease, compositions for diagnostics.

Four peptide sequences from Epstein-Barr virus were separately evaluated for binding to sera from patients with an autoimmune disease (Figures 3 and 7). All are found in the EBNA-1 (Epstein-Barr virus Nuclear Antigen-1) protein. Subjects with an autoimmune disease, systemic lupus erythematosus, tend to have higher levels of antibodies against PPPGRRP, GRGRGRGG and RGRGREK than do normal controls. On the other hand the glycine-alanine repeat sequence, GAGAGAGAGAGAGAGAGAGA, which, after infection by Epstein-Barr virus, is a major epitope in infectious mononucleosis and in normals (Rhodes, G. et al. J. Exp. Med. 165:1026-1040 (1987)) tends not to be bound by patients with lupus (Figure 7).

(SEQ ID NO: 1) (SEQ ID NO: 2) (SEQ ID NO: 3)  
 GA (SEQ ID NO: 7)

The overlapping octapeptides were constructed from the sequence of EBNA-1 (Epstein-Barr virus Nuclear Antigen-1) protein and preliminarily evaluated in four lupus patient sera and four controls. One control had no serologic evidence of having been exposed to Epstein-Barr virus, while the other three have anti-Epstein-Barr virus Viral Capsid Antigen IgG antibodies. All four of the patients had anti-Epstein-Barr virus Viral Capsid Antigen IgG antibodies.

The binding to octapeptides from EBNA-1 (Epstein-Barr virus Nuclear Antigen-1) protein shows dramatically different patterns between the Epstein-Barr virus exposed lupus patients and normal controls (Figure 8). These and other differences could become the basis for diagnostics that predict the risk of an autoimmune disease, that are associated with the presence of an autoimmune disease, and that are associated with particular clinical findings or manifestations. The structures bound by the lupus sera tested are listed in Table #.6

1



6  
 Table 5. Octapeptides from Epstein-Barr virus Nuclear Antigen-1 bound by the sera from two patients with systemic lupus erythematosus.

Sybil  
B9

Octapeptide Number	Sequence	Vaughan
30-48	GPQRRGGDNHGRGRGRGRGRGGGRPG	
58-70	GGSGSGPRHRDGVRRPQKRP	
72	RPQKRPS	
74-83	QKRPSICIGCKGTHGGTG	
88-93	GTGAGAGARGRG	
334	SGGRGRGG	e1*
347-351	RGSGGRRRGR	
368-376	RARGRGRGRGEKRPRS	e4*
388-394	SSSSGSFPRPPPG	
397-414	RPPPGRRPFFHPVGEADYFEYHQEG	
427	PDVPPGAI	
431	PGAIEQGPA	
445	GPSTGPRG	
452-453	GQGDGGRK	e14*
455-462	DGGRRKKGGWFGKHR	e14*
466-468	GKHRGQGSN	e11*
470	GQGSNPK	
475	NPKFENIA	
491	RSHVERTT	
508	VFVYGGSKT	
512-513	GSKTSLYNL	
542	GMAPGPGP	e12*
549	PQPGPLRE	
591	CNIRVTVC	
594-596	RVTVCSEDDG	
<del>607-608</del>	<del>PPWFPPMVEG</del>	<del>e10*</del>

These are taken from the data in Figures 8D and 8E. The peptides presented were bound in one of the sera by greater than 0.450 Absorbance Units. Sequences longer than eight amino acids represent neighboring octapeptides that exceed the 0.450 A.U.

threshold. Studies by John Vaughan and colleagues found some peptides of Epstein-Barr virus Nuclear Antigen-1 to be more antigenic in lupus patients than in controls (Petersen, et al, Arthritis Rheum. 33: 993-1000 (1990)). The octapeptides which share sequence with the peptides that were used in the Peterson study are identified with the peptide name used in their study (Table 6). Those peptides which they found to be differentially bound by lupus sera, relative to control sera, are identified with an \*.

Other differences in the antibody and cellular responses are expected to be important for the purpose of predicting the presence of (diagnosis) or subsequent risk of an autoimmune disease. Those presented above are examples of structures potentially useful for this purpose.

Assuming Epstein-Barr virus causes some autoimmune diseases, then differences in the immune response against this virus have the potential to predict risk of autoimmune disease and to be an aid to diagnosis and management of autoimmune disease.

Other structures defined by the proteins, glycoproteins nucleic acids, etc., will also be useful for this purpose. Such diagnostic tests can be based upon the relative presence of an antibody, cellular proliferation, molecular binding, cytokine production, skin reaction (erythema or induration), cell surface antigen or other measure of activation.

**Example 9. Use of a vaccine composition designed to induce a response to prevent or treat an autoimmune disease.**

Assuming Epstein-Barr virus causes autoimmune disease, then an effective vaccine which induces a protective response against Epstein-Barr virus has the potential to protect the host from the autoimmune disease. This is particularly true if the structure(s) which induce autoimmune disease is(are) altered or removed from the vaccine.

Once infected, this virus is latent and in most, perhaps virtually all, individuals the virus emerges from latency at a low level throughout the remainder of life. The viral infected

cells or the immune response required to suppress the virus have the potential to be extremely important in inducing or sustaining the autoimmune disease. Cells latently infected by Epstein-barr virus may also alter the <sup>immune</sup>~~immune~~ response. Consequently, treatments designed to suppress or eliminate Epstein-Barr virus have the potential to ameliorate the symptoms and tissue damage of the autoimmune disease.

Treatments expected to be useful against autoimmune disease include compositions that suppress the emergence of Epstein-Barr virus from latency. Also, agents for gene therapy directed against Epstein-Barr virus will be useful. Biologicals may also prove useful against Epstein-Barr virus by altering the intracellular environment, making it less hospitable to the virus by directly affecting the virus or by making the immune response against the virus less prone to support an autoimmune disease process.

Modification and variations of the methods and composition of the present invention will be obvious to those skilled in the art from the foregoing detailed description of the invention. Such modifications and variations are intended to come within the scope of the following claims.

Ins  
BIO